

Two-years surveillance of fungal contamination in three hospital departments in Campania Region

L. BRUNETTI, E. SANTORO, P. CAVALLO, G. BOCCIA, O. MOTTA, M. CAPUNZO

Department of Educational Science, Chair of Hygiene, Laboratory of Hygiene and Occupational Medicine, University of Salerno, Italy

Key words

Hospital • Environmental surveillance • Air and surfaces sampling • *Aspergillus* spp • Airborne fungal spores

Summary

A two-years (2003-2005) fungal environmental surveillance was carried out in three departments of a hospital in Campania region (Surgery, Intensive Care Unit, Obstetrics and Gynaecology). Four operating theatres rooms and their relative areas of service and support, 4 patient rooms of intensive care unit, 1 delivery room, 1 labour room and 1 nursery of Obstetrics and Gynaecology were checked. A total of 12,120 surfaces and 2,904 air samples were collected in 24 monthly determinations. A seasonal variation in the fungal development was observed, in particular the lowest level of air and surface fungi contamination was found in winter and autumn whereas it was higher in spring and summer. In this study 30 fungal species were identified and,

among these, the most frequent specie isolated was the *Aspergillus* spp. The results show an air contamination, expressed in percentage of positive determinations for *Aspergillus* spp, and the other fungi in the following percentages: Obstetrics and Gynaecology (25% and 33,3%); Intensive Care Unit (17% and 25%); Surgery (12.5% and 21%). For surfaces contamination it was found: Obstetrics and Gynaecology (67% and 75%); Intensive Care Unit (63% and 71%); Surgery (58.3% and 67%). This study shows that in the departments observed environmental fungi contamination is always present and therefore it would be necessary to apply environmental surveillance procedure and monitor the effectiveness.

Introduction

Airborne fungal spores have been widely recognized as major allergens capable of causing asthma and allergic rhinitis as well as other diseases. The environmental fungi contamination is a high risk factor for the generation of opportunist pathologies [1, 2] especially in the hospitals where the concentration of people highly exposed is elevated (e.g. those receiving cytotoxic therapy for haematological malignancies or undergoing organ or haematopoietic stem cell transplantation and post operate). The main fungal genera responsible for these infections are: *Aspergillus* spp, *Fusarium* spp, *Scedosporium* spp and *Mucorales* [3, 4]. However, any filamentous fungus can be potentially pathogen [5, 6]. The inhalation of airborne *Aspergillus* spores, either directly or through intermediate nasopharyngeal colonization, is a direct cause of pulmonary infection in immunocompromised patients [7, 8]. The *Aspergillus* spores are widespread, since they originate from decaying vegetation and are spread over by the air [7, 8]. In some studies, there are evidences that also houses [9-11] and schools [12-16] are particularly exposed to such a contamination.

Possible sources of airborne nosocomial infection in hospital include ventilation or air-conditioning systems, decaying organic material, dust, ornamental

plants, food, water and, particularly, building works in and nearby hospital [17, 18].

The aim of this study was an exploratory monitoring of possible fungi contamination in a hospital plexus, carried out by sampling several departments and their relative areas of service and support.

Materials and methods

ENVIRONMENTAL SAMPLING AND FREQUENCY

The study lasted two years (February 2003-March 2005) with a monthly frequency of determination. For details about the number of air and surfaces samples obtained at each determination see Table I. The three departments investigated are: Surgery, Intensive Care Unit, Obstetrics and Gynaecology.

The total number of samples for all determinations and for each department is reported in Table II.

ENVIRONMENTAL MONITORING

Environmental contamination was monthly examined (during two-years surveillance) by surface and air sampling, according to the protocol described in Table I. A total of 12,120 surfaces samples were taken, using contact plate RODAC (55 mm) containing Sabouraud Chlo-

Tab. I. Environmental sampling of air and surfaces in the departments and number of samples in each site for each determination. Air sampling: 1,000 L per each site. Surfaces were sampled by contact plate RODAC.

Department	Environment/ Environment Number	Air sample Number per Environment	Surface sample Number per Environment
Surgery	infirmary/1	3 samples random	10 samples on: hand basin, walls, windows, trolley, door, furniture, shower
	Sterilization room/2	5 samples random	20 samples on: hand basin, walls, windows, door, furniture equipment
	Pre operating room/3	3 samples random	15 samples on: hand basin, walls, windows, door
	Operating room/3	6 samples random	25 samples on hand basin, walls, windows, door, trolley
	Awakening room/2	6 samples random	30 samples on bed, walls, windows, door, trolley furniture
Intensive Care Unit	emergency room/1	3 samples random	20 samples on: hand basin, walls, windows, door, furniture equipments, trolleys
	corridor/3	6 samples random	10 samples on: walls, windows, door, equipment,
	Patient room/3	6 samples random	30 samples on hand basin, walls, windows, door, furniture equipment, trolley, bed
Obstetrics and Gynaecology	Sterilization room 1	5 samples random	15 samples on: hand basin, walls, windows, door, furniture equipments, trolleys
	Pre operating room/1	3 samples random	25 samples on: hand basin, walls, windows, door
	Operating room/1	6 samples random	30 samples on: hand basin, walls, windows, door, trolleys
	Delivery room/1	6 samples random	20 samples on: hand basin, walls, windows, door, equipments
	Labour room/1	5 samples random	20 samples on: hand basin, walls, windows, door, furniture equipments, beds
	Nursery/1	5 samples random	25 samples on: hand basin, walls, windows, door, equipments

ramphenicol agar, in order to evaluate whether fungi spores were present. The floor, which is supposed to be the area of major contamination, was excluded. A total of 2,904 air samples were collected (1,000 L sampled at 200 L/min) using a SAS mod. Super 180 apparatus (International pbi Spa Milano) loaded with a 55 mm diameter Petri dish, containing the same medium of contact plate RODAC. Petri dishes were incubated at 37 °C and examined after 48-72 hours, counting the number of colony growth. The fungi species observed were identified by their macroscopic aspect and by microscopic aspect of their spores after lacto phenol cotton blue staining.

Results

As reported in Table III the fungal contamination more frequently found was *Aspergillus* spp. The minimum average value in the air was 2.5 cfu/m³ in the Surgery and the maximum 3.8 cfu/m³ in the Obstetrics and Gynaecology department, whereas for the other species a minimum of 4.5 cfu/m³ was found in the Intensive Care Unit and a maximum average value of 5.1 cfu/m³ in the Obstetrics and Gynaecology.

The following species were identified: *Mucorales* spp, *Penicillium* spp, *Paecilomyces* spp, *Fusarium* spp, *Epicoccum* spp, whereas *Chaetium* spp and *Cephalosporium* spp were only occasionally found.

Tab. II. Total number of samples for all determinations and for each department.

Determination numbers = 24	Department							
	Surgery		Intensive Care Unit		Obstetrics and Gynaecology		Total samples	
	Air	Surf.	Air	Surf.	Air	Surf.	Air	Surf.
	1,248	5,520	936	3,360	720	3,240	2,904	12,120

With regard to the surfaces analyzed *Aspergillus* spp was found with a minimum value of 7.7 cfu/Petri dish in the Surgery and a maximum value of 9.2 cfu/Petri dish in the Obstetrics and Gynaecology. In the department of Obstetrics and Gynaecology *Aspergillus* contamination was localized particularly on the equipments, on the walls and windows, whereas in other departments it was found to be diffused on all the surfaces sampled. The other species identified, with a minimum value of 10.5 cfu/Petri dish in the Surgery and a maximum value of 14.3 cfu/Petri dish in the Obstetrics and Gynaecology, are the following: *Mucorales* spp, *Penicillium* spp, *Paecilomices* spp, *Fusarium* spp, *Epicoccum* spp, *Chaetium* spp and *Cephalosporium* spp, whose contamination was found to be diffused on all the surfaces investigated. *Paecilomices* spp, *Mucorales* spp and *Penicillium* spp, were prevalent on the tops of the furniture and over the equipments, compared with other species which, instead, were prevalent on the other surfaces. *Trichoderma* spp was only occasionally found on the headboards and *Candida* spp on the hand basins.

Discussion

The pulmonary infection as complication in hospitalized patients, is very frequent. This possibility is more dangerous in immunocompromised patients [2]. Normally there is no specific investigation about the causes of these infections, that very often are due to environmental fungi contamination.

In this work we performed a long period monitoring of hospital environments in a health care district of the

Campania region and analyzed the presence of fungi spores on the surfaces and in the air.

Air and surfaces monitoring shows a fungi contamination, that even if do not associated to infection cases of such an origin in this hospital plexus, to our knowledge, represents a high risk factor both for patients and for hospital employers. The lack of a filtering system capable of an high grade purification, together with non pressurized environment plays a crucial role in the fungal contamination.

Therefore a valid prevention of fungi infection risk would be the installation, in these environments, of adequate air purification systems and to monitor their efficiency so that spores fungi accumulation on the surfaces would be prevented [3, 19-21].

This study shows an high air fungi contamination in all the departments investigated, mostly in the Obstetrics and Gynaecology where we found the maximum frequency of positive determinations and the highest average values of both *Aspergillus*' spores and other fungi. The presence in the air of *Aspergillus*' spores in the concentrations observed is a high risk factor for Invasive Aspergillosis (IA) [1]. With regard to the surfaces contamination there is evidence of a higher percentage of positive determinations compared with the results of air determinations in all the departments investigated. This phenomenon can be connected with dust accumulation, that favours the deposit of pathogenic fungi spores [22], and which is probably due to a non frequent or insufficient environment cleaning [2]. Among the surfaces detected the ones that show major contamination are furniture tops and top surfaces of big equipments (refrigerators, sterilization apparatus, heaters,

Tab. III. Cumulative rates of fungi contamination of air and surfaces during the study period (2003-2005). For each department, the total number and percentage of positive determination during the study period are indicated in the upper row. Average and range of fungal contamination for *Aspergillus* and other fungi are given in cfu/m³ for air and in cfu/Petri dish for surfaces, in the lower row.

Department	Air 24 determination		Surfaces 24 determination	
	<i>Aspergillus</i>	Other Fungi	<i>Aspergillus</i>	Other Fungi
Surgery	3 12.5%	5 21%	14 58.3%	16 67%
	2.5 (2-9)	4.9 (3-15)	7.7 (3-20)	10.5 (8-25)
Intensive care unit	4 17%	6 25%	15 63%	17 71%
	3.4 (2-11)	4.5 (4-20)	8.8 (4-16)	12.1 (3-20)
Obstetrics and Gynaecology	6 25%	8 33.3%	16 67%	18 75%
	3.8 (1-10)	5.1 (4-18)	9.2 (4-35)	14.3 (5-40)

etc.) where dust accumulation is easier to occur [23, 24]. The species of fungi found on the surfaces are closely correlated with the ones found in the air; the department with major surfaces contamination being Obstetrics and Gynaecology.

In conclusion, in all the departments object of our study, an environmental fungi contamination was found, both in the air and on the surfaces, with small variation between them.

References

- [1] Alberti C, Bouakline A, Ribaud P, Lacroix C, Rousselot P, Leblanc T, Derouin F. *Relationship between environmental fungal contamination and the incidence of invasive aspergillosis in haematology patients*. J Hospital Infection 2001;48:198-206.
- [2] Heinemann S, Symoens F, Gordts B, Jannes H, Nolard N. *Environmental investigations and molecular typing of Aspergillus flavus during an outbreak of prostoperative infections*. J Hospital Infection 2004;57:149-55.
- [3] Diaz-Guerra TM, Mellado E, Cuenca-Estrella M, Gaztelurrutia L, Villate Navarro JJ, Rodriguez Tudela JL. *Genetic similarity among one Aspergillus Flavus strain isolated from a patient who underwent heart surgery and two environmental strains obtained from the operating room*. F Clin Microbiol 2000;38:2419-22.
- [4] Gosbell IB, Morris ML, Gallo JH. *Clinical, pathologic and epidemiologic features of infection with Scedosporium prolificans: four cases and review*. Clin Microbiol Infect 1999;5:672-86.
- [5] Guarro J, Nucci M, Akiti T, Gene J, Cano J, Barreiro MD, et al. *Phialemonium fungemia: two documented nosocomial cases*. F Clin Microbiol 1999;37:2493-7.
- [6] Hennequin C, Chouaki T, Pichon JC, Strunski V, Raccurt C. *Otitis externa due to Trichoderma longibrachiatum*. Eur F Clin Microbiol Infect Dis 2000;19:641-53.
- [7] Vandenberg MFQ, Verweij PE, Voss A. *Epidemiology of nosocomial fungal infections: Invasive Aspergillosis and the environment*. Diagn Microbiol infect Dis 1999;34:221-7.
- [8] Tablan OC, Anderson LJ, Arden NH, Breiman RF, Butler JC, McNeil MM. *Guideline for prevention of nosocomial pneumonia*. Respir Care 1994;39:1191-230.
- [9] Hagneaves M, Parappukkaran S, Morawska L, Hitchins J, He C, Dale G. *A pilot investigation into associations between indoor airborne fungal and non-biological particle concentrations in residential houses in Brisbane, Australia*. Sci Total Environ 2003;312:89-101.
- [10] Dybendal T, Elsayed S. *Dust from carpeted and smooth floors VI. Allergens in homes compared to those in schools in Norway*. Allergy 1994;49:210-6.
- [11] Gyntelberg F, Suadicani P. *Dust and the sick building syndrome*. Indoor Air 1994;4:223-38.
- [12] Daisey JM, Angell WJ, Apte MG. *Indoor air quality, ventilation and health symptoms in schools: an analysis of existing information*. Indoor Air 2003;13:53-64.
- [13] Levetin E, Shaughnessy R, Fisher E. *Indoor air quality in schools: Exposure to fungal allergens*. Aerobiologia 1995;11:27-34.
- [14] Dacarro C, Picco AM, Grisoli P, Rodolfi M. *Determination of aerial microbiological contamination in scholastic sports environments*. J Appl Microbiol 2003;95:904-12.
- [15] De-Wei L, Kendrick B. *Indoor aeromycota in relation to residential characteristics and allergic symptoms*. Mycopathologia 1995;131:149-57.
- [16] Smedje G, Norback D. *Irritants and Allergens at School in Relation to Furnishings and Cleaning*. Indoor Air 2001;11:127-33.
- [17] Parat S, Fricker-Hidalgo H, Perdrix A, Bemer D. *Airborne fungal contamination in air-conditioning systems: effect of filtering and humidifying devices*. Am Ind Hyg Assoc F 1996;57:996-1001.
- [18] Daniau C, Kaouffmann-Lacroix C, Castel O. *L'aerobioccontamination fongique en milieu hospitalier*. F Mycol Med 1998;8:139-46.
- [19] Cornet M, Levy V, Fleury L, Lortholary J, Barquins S, Coureul MH, et al. *Efficacy of prevention by high-efficiency particulate air filtration or laminar airflow against Aspergillus airborne contamination during Hospital renovation*. Infect Control Hosp Epidemiol 1999;20:508-13.
- [20] *Prevention du risqué Aspergillaire chez les patients immunodéprimés (Hématologie, transplantation)*. Hygiène 2000;8:305-504.
- [21] Derouin F. *Aspergillose invasive nosocomiale. Diagnostic, prévention et moyens de lutte intégrée en milieu hospitalier*. Bull Acad Natl Med 1996;180:859-70.
- [22] Leenders AC, van Belkum A, Behrendt M, Luijendijk A, Verbrugh HA. *Density and molecular epidemiology of Aspergillus in air and relationship to outbreaks of Aspergillus infection*. F Clin Microbiol 1999;37:1752-7.
- [23] Lebeau B, Pinel C, Grillot R, Ambroise-Thomas P. *Related Articles. Fungal and parasitic nosocomial infections: importance and limitations of disinfection methods*. Path Biol 1998;46:335-40.
- [24] Brocard-Lemort C. *Normes et recommandations en hygiène environnementale hospitalière*. Ann Biol Clin 2000;58:431-7.

■ Received on June 17, 2005. Accepted on November 3, 2005.

■ Acknowledgements: The authors of this study wish to thank Dr Ugo Ceravolo for his precious collaboration.

■ Correspondence: Dr Luigi Brunetti, Department of Educational Scienze, Chair of Hygiene, Laboratory of Hygiene and Occupational Medicine, via Ponte Don Melillo, 84084 Fisciano (SA), Italy. Tel. +39 089 962795 – Fax +39 089 962191 – E-mail: lbrunetti@unisa.it