

ORIGINAL ARTICLE

Environmental microbial contamination in dental setting: a local experience

M. GUIDA, F. GALLÉ*, V. DI ONOFRIO*, R.A. NASTRO, M. BATTISTA**, R. LIGUORI*, F. BATTISTA**, G. LIGUORI*
 Department of Structural and Functional Biology, University of Naples "Federico II"; * Chair of Hygiene and Epidemiology,
 Department of Studies of Institutions and Territorial Systems, University of Naples "Parthenope"; ** Complex Operating
 Unit of Odontostomatology and Maxillo-Facial Surgery, Pellegrini Hospital, Naples, Italy

Key words

Microbial contamination • Dental clinic • Environmental microbiological monitoring

Summary

Introduction. Patients and operators are exposed during dental practice to an infective risk, which derives especially from microorganisms suspended in aerosols. Environmental microbiological monitoring in dental settings represents a good instrument to detect critical situations.

Methods. In order to investigate environmental microbial contamination level in a local reality, we analyzed water, air and surfaces samples of a community-based dental facility by using protocol and threshold values proposed in a recent multicenter study carried out by the Italian Society of Hygiene, Preventive Medicine and Public Health (S.It.I.) working group "Hygiene in Dentistry". Microbial contamination was assessed in the same room for 4 non-consecutive weeks during all the five working days, before and at the end of the daily activity. Air was sampled also during clinical activity, through both active and passive sampling systems.

Results. Contamination of water showed a decrease during activities, while a decrease in air contamination was registered only at the end of the day. Passive sampling values resulted more often above threshold values adopted. At the same time, surfaces contamination increases at the end of the activity. It seems that in the dental clinic analyzed microbial buildup represents the higher critical element. No differences have been registered among the different days of the week.

Discussion. Our study highlights the need to improve disinfection procedures and air treatment systems in the considered environment. Microbiological monitoring could represent an important element to detect the presence of risk factors and to adopt control measures.

The full article is free available on www.jpmh.org

Introduction

Patients and staff of dental clinics are frequently at risk for infections. They are regularly exposed to water from aerosols generated during work, therefore the quality of dental unit water is of great importance, and Dental Unit Water Line (DUWL) contamination has become alarming [1]. The presence of opportunistic and pathogenic bacteria, such as *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Legionella pneumophila* in dental units water has been reported in several studies [2-7].

The presence of bacterial biofilms on the inside of DUWLs has been well documented and recognized as an undisputed source of contamination for water used in patient dental treatment [8, 9]. The risk of acquiring infections through Dental Unit Water Systems (DUWS) supplies is known to be not very uncommon. Often potential pathogenesis can spread through surgical procedures, local mucosal contact, ingestion and inhalation [10].

In particular, aerosols containing biological material (saliva, blood, and dental plaque) and microorganisms are produced by high-velocity rotating instruments and air-water sprays or ultrasonic scalers [11, 12]. Splatter and aerosols from dental procedures may infect health care personnel [13].

Smaller particles (<5 µm) can float in the air and have the potential to penetrate in the small passages of the lungs, while larger ones settle easily onto environmental surfaces which can become contaminated during patient care [12, 14-18].

Certain surfaces, especially clinical contact surfaces that are frequently touched (e.g. dental unit switches, light handles, drawer knobs), can act as reservoirs of microorganisms. When these surfaces are touched, microorganisms can be transferred to instruments, other environmental surfaces, or to the nose, mouth or eyes of health-care workers or patients [19].

Despite everything, data on microbial contamination in the dental clinic environment are not exhaustive [11, 12, 15, 16, 18-26].

Moreover, while the evaluation of water contamination is based on generally accepted and standardized sampling and processing protocols and well defined threshold values [27, 28], methods for air and surface sampling are still debated [28, 29] and no recommended levels of contamination for dental clinics have been established.

Since its founding, the working group "Hygiene in Dentistry" of the Italian Society of Hygiene, Preventive Medicine and Public Health (S.It.I.) analyzes the aspects

of microbial environmental contamination in dental clinics [30-33].

Recently, in order to establish a benchmark model for dental units microbial contamination, and considering the critical issues that have been observed in a previous pilot study, the working group carried out a wider multicentre study to assess water, air and surfaces microbial contamination using a standardized protocol [34, 35].

In this study, we analyzed the water, air and surfaces microbial contamination level of a community-based dental facility in the Hospital "Vecchio Pellegrini" of Naples by applying that protocol and by using threshold values proposed in the multicenter study.

The purposes of this study were to provide an accurate description of environmental microbial contamination in a local reality, and, secondly, to evaluate the results obtained on the basis of the benchmark proposed in the previous study.

Materials and methods

The study was carried out during the spring 2012. The environment analyzed was used for conservative and surgical practices. In order to evaluate weekly and daily modifications, microbial contamination of water, air and surfaces was assessed in the same room during the five working days of the week, before (T_0) and at the end of the daily activity (T_2), immediately after the last patient. In addition, air was sampled also during (T_1) clinical activity. Sampling was repeated for 4 non-consecutive weeks.

EVALUATION OF MICROBIAL WATER CONTAMINATION

Water from the tap and from dental unit water systems (DUWS) was sampled to assess the Total Viable Count (TVC) at 36°C and 22°C, and the presence of *Legionella* spp. and *P. aeruginosa*. For DUWS, the TVC was measured on every handpiece (1 ml of water from cup filler, air-water syringe, turbine, microengine, ablator), while *Legionella* spp. and *P. aeruginosa* were measured from a single sample made up collecting water from all the

five handpieces (200 mL from each handpiece for *Legionella* spp. and 50 mL from each handpiece for *P. aeruginosa*). All samples were collected in sterile bottles and immediately transported in a cool box (4-8°C) to the laboratory.

The Italian legislation n. 31 of February 2, 2001, based on the European Council Directive 98/83/EC on the quality of water intended for human consumption (1998), was used as the reference for TVC and *P. aeruginosa* detection [27, 36]. *Legionella* identification was performed according to the Italian guidelines for the prevention and control of legionellosis [37].

Water samples were processed as previously reported [34]. For TVC, Plate Count Agar plates were incubated at 22°C for 72 h and at 36°C for 48 h. Median values were compared with the threshold values established by both the European Council Directive 98/83/EC (20 cfu/mL at 36°C and 100 cfu/mL at 22°C) and CDC Guidelines (500 cfu/mL) [27, 28].

As for *P. aeruginosa* detection, Cetrimide agar incubated at 37°C for 48 h and API 20 NE Galleries (Bio Mérieux Srl, Charbonnières-les-Bains, France) for biochemical confirmation were used. Absence of bacteria in 250 mL of sampled water was considered as threshold value [27].

To determine the presence of *Legionella* spp., we used glycine-vancomycin-polymyxin-cyclohexamide (GVPC) agar medium. Suspect colonies were subcultured on charcoal-yeast-extract (CYE) agar and buffered-charcoal-yeast-extract (BCYE) agar. Serological identification was performed by latex test, and by a monoclonal antibodies trial (DenkaSeiken Co. LTD, Tokyo, Japan). Regarding the presence of *Legionella* spp., a threshold value of 1,000 cfu/L was used [37] (Tab. I).

EVALUATION OF MICROBIAL AIR CONTAMINATION

Microbial air contamination was evaluated through both active and passive sampling [38]. Active sampling was performed using the Surface Air System (SAS) sampler (International PBI, Milan, Italy), with a flow rate of 180 liters per minute (L/min) and a suction volume of 500 L.

Tab. I. References adopted for contamination threshold values.

Reference	Water			Air		Surfaces	
	TVC 22°C cfu/mL	TVC 36°C cfu/mL	<i>Pseudomonas aeruginosa</i> (cfu/250 mL)	<i>Legionella</i> spp. (cfu/L)	Active Sampling (cfu/m ³)	Passive Sampling (cfu/plate 9cmØ/h)	Countertop Dental unit switches
European Council Directive	100	20	0				
CDC Guidelines	500	500					
Italian Guidelines for the prevention and control of legionellosis				1,000			
Pasquarella et al., Sci Tot Environ 2012					Target 166 Alert 278	Target 27.5 Alert 43.7	0.64 0.63

The number of colony forming units was adjusted using the conversion table provided by the manufacturer and was expressed in colony forming units per cubic meter (cfu/m³).

Passive sampling was performed to determine the number of cfu grown on a Petri dish of a 9 cm diameter placed for 1 hour on a surface (Index of Microbial Air contamination, IMA) [29].

Both samplers were placed in the monitored room about 1 m above the floor and about 1 m away from dental unit, walls and any other obstacle.

Total Viable Count (TVC) was performed using Tryptone Soya Agar plates incubated at 36°C for 48 h.

Presence of *Legionella* spp. was evaluated only through active sampling, using RODAC plates containing glycine–vancomycin–polymyxin–cyclohexamide (GVPC) agar.

Target and alert values proposed by Pasquarella et al. have been used as benchmark to evaluate the contamination level of the air (Tab. I) [35].

EVALUATION OF SURFACE MICROBIAL CONTAMINATION

Microbial contamination of the countertop serving the monitored dental unit and that of dental unit switches surfaces were evaluated. A RODAC plate, 55 mm in diameter, was pressed on the surface to be tested, and then incubated at 36°C for 48 h. The results were expressed as cfu/cm².

Threshold values reported by Pasquarella et al. have been used to evaluate the contamination level of the surfaces (Tab. I) [35].

STATISTICAL ANALYSIS

Descriptive statistical analysis was performed to provide median, minimum-maximum values range, and mean and standard deviation.

Parametric or nonparametric tests were applied on the basis of data distribution.

The Wilcoxon signed-rank test was used to analyze differences in microbial water contamination median values between T₀ and T₂, and those in microbial surface contamination median values between T₀ and T₂. Differences in microbial air contamination mean values

between T₀ and T₁, and between T₁ and T₂ were evaluated by Student's t-test for paired samples. Differences among the microbial contamination recorded on different days of the week were evaluated by Friedman two-way Analysis of Variance by Ranks; possible differences among the results from Monday and Friday (the first and the last day of the weekly activity respectively) were also analyzed by using the Wilcoxon signed-rank test. Possible interference between the presence of *P. aeruginosa* and *Legionella* spp. was tested by one-tailed Fisher's exact test. A p value below 0.05 was considered statistically significant.

Results

Median values and ranges, together with mean and standard deviation for water contamination results are reported in Tables II and III. The number and percentage of tap and DUWS water samples above the threshold values were also reported.

As for tap water TVC, both median values at 36°C and at 22°C decreased from T₀ to T₂, but the reduction was statistically significant only for mesophilic counts (p = 0.005). This is confirmed by the decreasing number of samples which exceeded the limits established by the European Council Directive, registered only for those kept at 36°C (from 50% at T₀ to 30% at T₂). Both mesophilic and psychophilic contamination of water samples from DUWS showed a significant decrease (p = 0.00) from the start (T₀) to the end (T₂) of clinical practice. In total, *P. aeruginosa* was detected in 5 (12.5%) tap water samples and in 20 (50%) samples from DUWS, while *Legionella* spp. exceeded the limit value in 8 (20%) tap water samples and in 6 (15%) samples collected from DUWLs (data not shown). No statistically significant differences were found in *P. aeruginosa* contamination of both tap and DUWLs water (p = 0.27 and p = 0.16 respectively) collected before and at the end of the dental practice. The same results were registered for *Legionella* spp. contamination (p = 0.71 and p = 0.10 respectively). With one only exception, contamination by *P. aeruginosa* or *Legionella* spp. registered before the start of the activities at one sampling point was confirmed at the same

Tab. II. Microbial contamination values of water samples from tap water before (T₀) and after (T₂) clinical practice.

	T ₀				T ₂			
	TVC 22°C cfu/mL	TVC 36°C cfu/mL	<i>Legionella</i> cfu/L	<i>Pseudomonas</i> cfu/250mL	TVC 22°C cfu/mL	TVC 36°C cfu/mL	<i>Legionella</i> cfu/L	<i>Pseudomonas</i> cfu/250mL
Median	16.50	32.50	0	0	6	3.50	0	0
Range	0-9,000	0-39,000	0-3,700	0-8	0-300,000	0-300,000	0-5,300	0-6
Mean (SD)	1,140 (2,536.8)	2,937.5 (8,818.5)	420 (961.68)	0.6 (1.8)	16,118.1 (66,861.6)	15,108.75 (67,056.8)	540 (1,327.6)	0.45 (1.4)
N. (%) of samples above threshold value*	5 (25)	10 (50)	4 (20)	3 (15)	5 (25)	6 (30)	4 (20)	2 (10)

* Council Directive 98/83/EC 3 November 1998 on the quality of water intended for human consumption.

° Guidelines for the prevention and control of Legionellosis, 2000.

Tab. III. Microbial contamination values of water samples from dental unit water system before (T_0) and after (T_2) clinical practice.

	T_0				T_2			
	TVC 22°C cfu/mL	TVC 36°C cfu/mL	<i>Legionella</i> cfu/L	<i>Pseudomonas</i> cfu/250mL	TVC 22°C cfu/mL	TVC 36°C cfu/mL	<i>Legionella</i> cfu/L	<i>Pseudomonas</i> cfu/250mL
Median	61,000	23,500	0	1	9,000	5,000	0	0.5
Range	0-360,000	0-300,000	0-2,200	0-102	0-300,000	0-300,000	0-3,100	0-110
Mean (SD)	117,863.8 (1.2)	99,487.4 (121,762.3)	270 (675.9)	19 (34.6)	68,456.1 (1.1)	53,967.3 (98,868.4)	420 (1,032.4)	9.3 (24.3)
N. (%) of samples above threshold value*□°	88 (88) 87 (87)	93 (93) 83 (83)	3 (15)	10 (50)	78 (78) 72 (72)	86 (86) 68 (68)	3 (15)	10 (50)

* Council Directive 98/83/EC 3 November 1998 on the quality of water intended for human consumption.

□ CDC Guidelines for Environmental Infection Control in Health-Care Facilities, 2003.

° Guidelines for the prevention and control of Legionellosis, 2000.

Tab. IV. Prevalence of *Legionella* spp. and *Pseudomonas aeruginosa* in samples from dental unit water systems.

<i>Legionella</i> spp.	<i>Pseudomonas aeruginosa</i>		
	Negative n. (%)	Positive n. (%)	Total n. (%)
Negative	20 (50)	14 (35)	34 (85)
Positive	0 (0)	6 (15)	6 (15)
Total n. (%)	20 (50)	20 (50)	40

One-tailed Fisher's exact test $p = 0.01$

point at the end of the work. Table IV shows the relation between *Legionella* spp. and *P. aeruginosa* found in the DUWS. *Legionella* spp. was found more frequently when *P. aeruginosa* was present ($p = 0.01$). Six samples (15%) showed both *Legionella* spp. and *P. aeruginosa* presence.

Regarding air samples, Table V shows as median values of microbial contamination decreased only at the end of the clinical practice (T_2), without notable variation between T_0 and T_1 . The reduction was statistically significant only for passive sampling (33.5 to 15 cfu/plate, $p=0.001$) and not for active sampling (78 to 55.5 cfu/m³, $p = 0.736$). This is also confirmed by the number of samples above limit values adopted; only in a few cases active sampling gave samples out of the benchmarks. *Legionella* spp. was never detected.

Microbial surface contamination showed an increase between T_0 and T_2 both for countertops and dental unit switches, but this variation was not statistically significant ($p = 0.611$ and $p = 0.084$ respectively) (Tab. VI). Samples from dental unit switches exceeded threshold values more often at the end than at the start of the activities.

Neither water samples, nor air and surfaces samples, showed statistically significant differences in microbial contamination levels among the different days of the week. The comparison among results obtained on Mondays and Fridays did not give any significant difference, so for water, as for air and surfaces (data not shown).

Discussion

This study has been realized on the basis of the previous experiences of the SItI Working Group "Hygiene in Dentistry". In particular, monitoring procedures and limit values are the same adopted in the last multicenter studies carried out in six dental clinics of Italy [34, 35]. The results of this study support the importance of analyzing the environmental microbial contamination in dental settings, in order to control infective risks for patients and staff.

As previously reported [30, 34, 35], contamination of water from taps as for DUWS, which is probably consequent to night stagnation, showed a decrease during clinical activity. Contamination by *P. aeruginosa* was registered in a high number of samples (50%) from DUWS, while the contamination of tap water was always lower. In contrast to what registered in previous experiences, the presence of *Legionella* spp. was not favored by the lack of *P. aeruginosa*. On the contrary, they were both present in the 15% of the samples, and this probably demonstrates the presence of risk factors which could favor the development of critical situations.

The air contamination did not show increases during activity, while a decrease was registered at the end of the day, both for active and passive sampling. This could be explained with the presence of other factors independent by clinical activity, which can influence the quality of the air. Probably the status of air conditioning systems, which were often turned off at the end of the daily activity, could have an important role in determining these results.

Tab. V. Microbial air contamination values assessed by active sampling (cfu/m³) and passive sampling (IMA) before (T₀), during (T₁) and after (T₂) clinical practice.

	Active sampling (cfu/m ³)			Passive sampling (cfu/plate 9cmØ/h)		
	T ₀	T ₁	T ₂	T ₀	T ₁	T ₂
Median	75	78	55.5	36	33.5	15
Range	14-190	9-163	17-246	10-101	9-141	7-45
Mean (SD)	86.6 (50.6)	82.3 (48.3)	86.5 (64.8)	38.3 (21)	39.6 (28.2)	20 (11.5)
N. (%) of samples above target value*	2 (10)	0	3 (15)	13 (65)	13 (65)	5 (25)
N. (%) of samples above alert value*	0	0	0	7 (35)	6 (30)	1 (5)

* Benchmark proposed for dental clinics by Pasquarella et al. ³⁵.

Tab. VI. Microbial contamination values on countertop and switches before (T₀) and after (T₂) clinical practice.

	Countertop (cfu/cm ²)		Dental unit switches (cfu/cm ²)	
	T ₀	T ₂	T ₀	T ₂
Median	3,000	4,200	4,000	7,600
Range	0.08-2.44	0.04-2.16	0.04-2.64	0.12-10.40
Mean (SD)	0.47 (0.52)	0.57 (0.48)	0.61 (0.67)	1.5 (2.3)
N. (%) of samples above threshold value*	5 (25)	4 (20)	4 (20)	11 (55)

* Benchmark proposed for dental clinics by Pasquarella et al. ³⁵.

IMA values resulted more often above both target and alert values proposed by Pasquarella et al. [35]. Since passive sampling has the role to evidence risks deriving from particles fallout, while active sampling has to reveal those deriving from aerosols diffusion [39], it seems that in the analyzed dental clinic settled particles could represent the higher critical element.

This was also demonstrated by surfaces biocontamination, which increases at the end of the activity. This surely represents the effect of working activities, but probably it could also derive from the fallout of airborne particles. Here, as in other experiences of our Working Group, Rodac plates represented the best choice among the surface sampling systems, both for accuracy and user-friendliness [34, 35, 40].

Surprisingly, no differences between the contamination levels at the start (on Monday) and at the end (on Friday) of weekly clinical activity have been registered.

Evidently, facility closing along the weekend did not increase the effects of water stagnation and microbial particles fallout insomuch as the levels of microbial contamination show a significant modification. Regarding air contamination, the switching off of malfunctioning air conditioners, as above suggested, could have contributed to this result.

Conclusions

In conclusion, data reported highlight the need to improve disinfection procedures and air treatment systems in the considered environment, so during as at the end of the activities. Furthermore, our study underlines that microbial monitoring could represent an important element to detect alert values which indicate the presence of risk factors and require the adoption of control measures.

References

- [1] Szymańska J, Sitkowska J, Dutkiewicz J. *Microbial contamination of dental unit waterlines*. Ann Agric Environ Med 2008;15:173-9.
- [2] Al-Hiyasat AS, Ma'ayeh SY, Hindiye MY, et al. *The presence of Pseudomonas aeruginosa in the dental unit waterline systems of teaching clinics*. Int J Dent Hyg 2007;5:36-44.
- [3] Pankhurst CL, Philpott-Howard JN. *The microbiological quality of water in dental chair units*. J Hosp Infect 1993;23:167-74.
- [4] Atlas RM, William JF, Huntington MK. *Legionella contamination in dental unit water*. Appl Environ Microbiol 1995;61:1208-13.
- [5] Challacombe SJ, Fernandes LL. *Detecting Legionella pneumophila in water systems: a comparison of various dental units*. J Am Dent Assoc 1995;126:603-8.
- [6] Barbeau J, Tanguay R, Faucher E, et al. *Multiparametric analysis of waterline contamination in dental units*. Appl Environ Microbiol 1996;62:3954-9.
- [7] Stampi S, Zanetti F, De Luca G, et al. *Effect of water softening and heating on microbial contamination of dental unit systems*. Zentralbl Hyg Umweltmed 1996;198:522-30.
- [8] Porteous N, Luo J, Herrera M, et al. *Growth and identification of bacteria in N-halamine dental unit waterline tubing using an ultrapure water source*. Int J Microbiol 2011;767314.
- [9] Donlan RM, Costerton JW. *Biofilms: survival mechanisms of clinically relevant microorganisms*. Clin Microbiol Rev 2002;15:167-93.

- [10] Fiehn NE, Larsen T. *The effect of drying dental unit waterline biofilms on the bacterial load of dental unit water*. Int Dent J 2002;52:251-4.
- [11] Cristina ML, Spagnolo AM, Sartini M, et al. *Evaluation of the risk of infection through exposure to aerosols and spatters in dentistry*. Am J Infect Control 2008;36:304-7.
- [12] Prospero E, Savini S, Annino I. *Microbial aerosol contamination of dental healthcare workers' faces and other surfaces in dental practice*. Infect Control Hosp Epidemiol 2003;24:139-41.
- [13] Wirthlin MR, Marshall GW Jr, Rowland RW. *Formation and decontamination of biofilms in dental unit waterlines*. J Periodontol 2003;74:1595-609.
- [14] Decraene V, Ready D, Pratten J, et al. *Air-borne microbial contamination of surfaces in a UK dental clinic*. J Gen Appl Microbiol 2008;54:195-203.
- [15] Harrel SK, Molinari J. *Aerosols and splatter in dentistry: a brief review of the literature and infection control implications*. J Am Dent Assoc 2004;135:429-37.
- [16] Rautemaa R, Nordberg A, Wuolijoki-Saaristo K, et al. *Bacterial aerosols in dental practice — a potential hospital infection problem?* J Hosp Infect 2006;64:76-81.
- [17] Szymanska J. *Dental bioaerosols as an occupational hazard in dentist's workplace*. Ann Agric Environ Med 2007;14:203-7.
- [18] Timmerman MF, Menso L, Steinfert J, et al. *Atmospheric contamination during ultrasonic scaling*. J Clin Periodontol 2004;3:458-62.
- [19] Centers for Disease Control and Prevention. *Guidelines for Infection Control in Dental Health-Care Setting, 2003a*. Available at <http://www.cdc.gov/oralhealth/InfectionControl/guidelines/index.htm>. Accessed June 29, 2011.
- [20] Coleman DC, O'Donnell MJ, Shore AC, et al. *Biofilm problems in dental unit water systems and its practical control*. J Appl Microbiol 2009;146:1-14.
- [21] Dutil S, Veillette M, Mériaux A, et al. *Aerosolization of mycobacteria and legionellae during dental treatment: low exposure despite dental unit contamination*. Environ Microbiol 2007;9:2836-43.
- [22] Kumar S, Atray D, Paiwal D, et al. *Dental unit waterlines: source of contamination and cross-infection*. J Hosp Infect 2010;74:99-111.
- [23] Pankhurst CL, Coulter WA. *Do contaminated dental unit waterlines pose a risk of infection?* J Dent 2007;35:712-20.
- [24] Pederson ED, Stone ME, Ragain Jr JC, et al. *Waterline biofilm and the dental treatment facility: a review*. Gen Dent 2002;50:190-5.
- [25] Rowland BM. *Bacterial contamination of dental unit waterlines: what is your dentist spraying into your mouth?* Clin Microbiol 2003;25:73-7.
- [26] Singh T, Coogan MM. *Isolation of pathogenic Legionella species and Legionella-laden amoebae in dental unit waterlines*. J Hosp Infect 2005;61:257-62.
- [27] Council Directive 98/83/EC of 3 November 1998 on the quality of water intended for human consumption: Official Journal of the European Community L330 1998:32-54.
- [28] Centers for Disease Control and Prevention. *Guidelines for Environmental Infection Control in Health-Care Facilities, 2003b*. Available at http://www.cdc.gov/hicpac/pdf/guidelines/eic_in_HCF_03.pdf. Accessed June 29, 2011.
- [29] Pasquarella C, Albertini R, Dall'Aglia P, et al. *Il campionamento microbiologico dell'aria: lo stato dell'arte*. Ig San Pubbl 2008;64:79-120.
- [30] Castiglia P, Liguori G, Montagna MT, et al. *Italian multicenter study on infection hazards during dental practice: Control of environmental microbial contamination in public dental surgeries*. BMC Public Health 2008;8:187-93.
- [31] Napoli C, Tatò D, Iatta R, et al. *Assessment of occupational risk of Legionella spp infection among dental healthcare personnel*. Ig Sanità Pubbl 2007;6:683-9.
- [32] Napoli C, Iatta R, Fasano F, et al. *Variable bacterial load of Legionella spp. in a hospital water system*. Sci Total Environ 2009;408:242-4.
- [33] Montagna MT, Tatò D, Napoli C, et al. *Pilot study on the presence of Legionella spp. in 6 Italian cities' dental units*. Ann Ig 2006;18:297-303.
- [34] Pasquarella C, Veronesi L, Castiglia P, et al. *Italian multicentre study on microbial environmental contamination in dental clinics: a pilot study*. Sci Total Environ 2010;408:4045-51.
- [35] Pasquarella C, Veronesi L, Napoli C, et al. *Microbial environmental contamination in Italian dental clinics: A multicenter study yielding recommendations for standardized sampling methods and threshold values*. Sci Total Environ 2012;420:289-99.
- [36] Decreto Legislativo 2 febbraio 2001, n. 31: "Attuazione della direttiva 98/83/CE relative alla qualità delle acque destinate al consumo umano". Gazzetta Ufficiale n. 52 del 3.3.2001, suppl. Ord. n. 41.
- [37] *Linee guida italiane per la prevenzione e il controllo della legionellosi*. Gazzetta Ufficiale della Repubblica Italiana n. 103 del 5.5.2000. Available at <http://legionellaonline.it>. Accessed June 29, 2011.
- [38] ISO 14698-1. *Cleanrooms and associated controlled environments – Biocontamination control. Part 1: General principles and methods*. 2003.
- [39] Liguori G, Spagnoli G, Agozzino E, et al. *Biological risk in the operating room: microbiological monitoring of the environment and analysis of the associated variables*. Ann Ig 2005;17:385-98.
- [40] Liguori G, Gombos F, Marinelli A, et al. *Microbial environmental monitoring in the dental surgery room*. Ann Ig 2003;15:123-33.

■ Received on September 8, 2012. Accepted on October 8, 2012.

■ Correspondence: Giorgio Liguori, Chair of Hygiene and Epidemiology, Department of Studies of Institutions and Territorial Systems, University of Naples "Parthenope", via Medina 40, 80133 Naples, Italy - Tel./Fax + 39 081 5474790 - E-mail: giorgio.liguori@uniparthenope.it