

Sociodemographic determinants of fruit and vegetable consumption in Indonesia: insights from the 2023 Indonesian Health Survey

AZIMATUL AMINI¹, FITRINA MAHARDANI KUSUMANINGRUM², DIGNA NIKEN PURWANINGRUM³, ABDUL WAHAB³, ADITYA LIA RAMADONA²

¹ Master of Public Health Program, Faculty of Medicine, Public Health, and Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia;

² Department of Health Behaviour, Environment, and Social Medicine, Faculty of Medicine, Public Health, and Nursing,

Universitas Gadjah Mada, Yogyakarta, Indonesia; ³ Department of Biostatistics, Epidemiology, and Population Health, Faculty of Medicine, Public Health, and Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia

Keywords

Dietary patterns • Health inequities • Health surveys • Noncommunicable diseases • Population health

Summary

Introduction. Inadequate fruit and vegetable consumption is a recognized risk factor for non-communicable diseases. This study aimed to identify the sociodemographic determinants of fruit and vegetable consumption in Indonesia, which would be fundamental to developing targeted public health interventions for non-communicable diseases.

Methods. This cross-sectional study utilized secondary data from the 2023 Indonesian Health Survey. We analyzed the data from 595,303 adults aged 19 years or older, comprising variables of fruit and vegetable consumption and sociodemographic determinants. A logistic regression analysis is conducted to examine associations between sociodemographic factors and fruit and vegetable consumption.

Results. A significant majority of Indonesians (97.1%) fail to meet the recommended intake of fruits and vegetables. Better fruit and vegetable consumption patterns were observed

among females ($aOR=1.31$; 95% $CI=1.24-1.38$), individuals with higher education ($aOR=1.47$; 95% $CI=1.23-1.75$), those employed ($aOR=1.13$; 95% $CI=1.06-1.20$), married ($aOR=1.25$; 95% $CI=1.16-1.34$), from lower economic status groups, and residents of Nusa Tenggara ($aOR=1.32$; 95% $CI=1.13-1.53$), Maluku ($aOR=1.54$; 95% $CI=1.16-2.06$), Papua ($aOR=1.95$; 95% $CI=1.46-2.60$), and rural areas ($aOR=1.12$; 95% $CI=1.02-1.23$).

Conclusion. Indonesians consume very few fruits and vegetables, and this low consumption isn't uniform. We see significant differences based on factors like gender, education level, occupation, marital status, socioeconomic status, and place of residence. These findings underscore the urgent need for targeted nutritional interventions and the formulation of context-specific public health policies to promote healthier dietary behaviors.

Introduction

In 2016, non-communicable diseases (NCDs) accounted for 73% of all deaths in Indonesia. The mortality breakdown consisted of 35% from cardiovascular diseases, 12% from cancer, 6% from diabetes mellitus, and 15% from other NCDs [1]. According to the national health survey Indonesian Basic Health Research (Riskesdas), the prevalence of NCDs in Indonesia has increased from 2013 to 2018 [2]. Prevention of risk factors contributing to NCDs must be prioritized in national health strategies to reduce the future prevalence of these diseases [3].

One of the primary risk factors for NCDs is inadequate consumption of fruits and vegetables (2). NCDs such as obesity are caused by unhealthy diets [4]. Changing unhealthy diets to healthy ones can reduce NCDs cases and help people achieve a healthier lifestyle [5]. Regular intake of fruits and vegetables can help reduce the risk of cardiovascular diseases, stroke, and hypertension; prevent certain types of cancer, such as breast, colon,

and lung cancer; and protect against neurodegenerative diseases like Alzheimer's and Parkinson's [6]. While these fruits and vegetables-NCDs relationship is established globally, their sociodemographic determinants in Indonesia remain underexplored despite rising NCDs [2] and current low consumption.

Kucuk (2023), in an analysis of the 2019 Turkish Health Survey, reported that fruit and vegetable consumption was positively associated with multiple sociodemographic and lifestyle variables - such as marital status, educational attainment, active behaviors (cycling and walking), adequate rest, higher income, place of residence, age, and total duration of physical activity [7]. While these insights underscore the multifaceted drivers of diet in Turkey, Indonesia's distinct cultural, economic, and geographic diversity may yield different patterns. Context-specific research is therefore indispensable for uncovering Indonesia's unique sociodemographic determinants of fruit and vegetable intake.

As non-communicable diseases continue to rise in Indonesia, and dietary habits shift alongside rapid

urbanization and income growth, the determinants of fruit and vegetable consumption remain underexplored. Addressing this gap is critical for developing well-targeted public health strategies. Accordingly, the primary objective of this study is to analyze the association between sociodemographic characteristics and fruit and vegetable consumption in Indonesia, using data from the 2023 Indonesian Health Survey (IHS). Specifically, we (1) characterized national patterns of fruit and vegetable consumption and (2) examined how these patterns vary across key sociodemographic subgroups, thereby informing targeted public health interventions.

Methods

STUDY DESIGN

This study employed a quantitative approach with a cross-sectional design, utilizing secondary data from the 2023 IHS. The IHS – known in Indonesian as Survei Kesehatan Indonesia (SKI) – is a nationally representative health survey conducted by the Indonesian Ministry of Health [8]. The 2023 IHS was selected for this study because it provides the most recent and comprehensive data on individual health status, sociodemographic characteristics, and regional variations. Data collection for the IHS was carried out between August and September 2023.

PARTICIPANTS AND DATA COLLECTION

The sample for this study consists of individuals aged 19 years and older, drawn from the 2023 IHS dataset. The independent variables include age, gender, education level, employment status, economic status, marital status, place of residence, and area classification. In the IHS dataset, economic status is assessed based on an index of household ownership of durable goods, which is then categorized into five quintiles. The dependent variable in this study is fruit and vegetable consumption. Data on fruit and vegetable consumption were collected using the STEPwise instrument developed by the World Health Organization (WHO) [8]. This study utilized the individual questionnaire from the 2023 IHS, with instruments sourced from both individual and household questionnaires. These were obtained from the official website of the Ministry of Health of the Republic of Indonesia (badankebijakan.kemkes.go.id/hasil-ski-2023/).

The sample included respondents who met the inclusion criteria, which required complete data on all relevant variables. Respondents were categorized into two groups based on their dietary behavior. Those who consumed an adequate amount of fruits and vegetables during the past week were classified as the “healthy group,” while those who did not were classified as the “less healthy group.” Adequate consumption was defined per WHO-STEPS as the intake of at least five servings of fruits and/or vegetables per day, for seven consecutive days [7, 8]. This WHO-STEPS-based cutoff exceeds Indonesia’s

Isi Piringku guideline (~3-4 servings/day across three meals) but enables international comparability.

STATISTICAL ANALYSIS

Initial dataset configurations were performed according to the guidelines outlined in the 2023 IHS Data User Manual. This setup included the application of sampling weights, as the IHS 2023 employed a complex sampling design, ensuring that the analytical results are nationally representative. Missing data were identified in the economic status variable ($n=46$). Given the small proportion relative to the total sample size ($n=593,303$), these cases were excluded from the analysis. Following data cleaning, variables were categorized according to the study’s inclusion criteria.

Descriptive statistical analysis was conducted to summarize the data. Frequencies and percentages were used to describe both independent and dependent variables, with univariate results presented in tabular format. Bivariate analysis was performed using the chi-square test and odds ratios (ORs). The chi-square test assessed the significance of associations between each sociodemographic variable and fruit and vegetable consumption, while ORs measured the strength of these associations. A significance level of $p<0.05$ was used for all bivariate tests.

Multivariable analysis was conducted using logistic regression. Only independent variables that demonstrated statistically significant associations in the bivariate analysis were included in the regression model. This approach was intended to minimize the influence of non-significant variables on the model’s outcomes.

Results

As presented in Table I, only 2.9% of respondents met the criteria for a healthy fruit and vegetable consumption pattern. The majority of participants (83.6%) were within the 19-59 age range. The sample was slightly skewed toward $\geq$ female respondents, with most individuals reporting marital status as married and employment status as employed. Educational attainment was predominantly at the primary level, representing 50.2% of the sample. Economic status was distributed relatively evenly across quintiles. Geographically, the largest proportions of respondents resided in Java (28.5%) and Sumatra (31.0%), with smaller distributions across Sulawesi, Kalimantan, Bali, Nusa Tenggara, Maluku, and Papua. Bivariate analysis (Tab. II) indicated statistically significant associations between fruit and vegetable consumption patterns and several sociodemographic variables, including gender, educational attainment, employment status, economic status, marital status, place of residence, and area classification. Age did not demonstrate a significant association and was therefore excluded from subsequent multivariable analysis.

Multivariable logistic regression analysis revealed that gender, education level, employment status, economic status, marital status, and regional residence remained

Tab. I. Description of respondents based on diet and sociodemographic characteristics.

Variables	N	%
N	593,303	100.0
Dietary pattern category		
Unhealthy dietary pattern	576,119	97.1*
Healthy dietary pattern	17,184	2.9
Age		
19-59 years	495,964	83.6
≥60 years	97,339	16.4
Gender		
Male	268,591	45.3
Female	324,712	54.7
Education level		
No formal education	30,188	5.1
Primary education	298,020	50.2
Secondary education	192,709	32.5
Higher education	72,386	12.2
Employment status		
Unemployed	196,292	33.1
Employed	397,011	66.9
Economic status		
Lowest	112,041	20.0
Lower-middle	113,283	20.2
Middle	110,865	19.8
Upper-middle	111,938	20.0
Highest	112,030	20.0
Marital status		
Unmarried	124,933	21.1
Married	468,370	78.9
Place of residence		
Jawa	168,918	28.5
Sumatra	183,951	31.0
Kalimantan	56,103	9.5
Sulawesi	88,832	15.0
Bali	13,730	2.3
Nusa Tenggara	36,100	6.1
Maluku	20,283	3.4
Papua	25,386	4.3
Area classification		
Urban	317,614	53.5
Rural	275,689	46.5

Source: IHS 2023. * Based on the WHO-STEPPS definition (≥5 servings of fruit and/or vegetables daily for 7 consecutive days). Using the less stringent Isi Piringku benchmark (~3–4 servings/day) would likely result in a lower estimated prevalence of inadequate intake.

significantly associated with healthy dietary patterns after adjusting for potential confounders. Female respondents were 1.31 times more likely to exhibit healthy fruit and vegetable consumption compared to male respondents (95% CI: 1.24–1.38). Higher education was positively associated with healthy dietary behavior, with respondents possessing higher education being 1.47 times more likely to meet recommended intake levels compared to those with no formal education (95% CI: 1.23–1.75). Employment status also showed a positive association, with employed individuals being 1.13 times more likely to

consume fruits and vegetables adequately than their unemployed counterparts (95% CI: 1.06–1.20).

Contrary to expectations, higher economic status was inversely associated with healthy fruit and vegetable consumption. Married individuals were 1.25 times more likely to adhere to healthy dietary patterns compared to unmarried individuals (95% CI: 1.16–1.34). Regional analysis indicated that respondents residing in Sulawesi, Nusa Tenggara, Maluku, and Papua were more likely to consume fruits and vegetables adequately compared to those in Java. In contrast, respondents from Sumatra, Kalimantan, and Bali exhibited lower odds of healthy consumption. Additionally, rural residence was associated with a higher likelihood of healthy dietary behavior compared to urban residence (aOR=1.12).

Discussion

The present analysis found that 97.1% of respondents were classified as having inadequate fruit and vegetable intake under the WHO-STEPPS definition (≥5 servings/day for 7 consecutive days). This high prevalence reflects the stringency of the operational cutoff, capturing shortfalls relative to an international benchmark, rather than implying an absence of fruit and vegetable consumption. Consequently, the outcome distribution is highly imbalanced (2.9% “healthy”), which is expected when applying a strict criterion in a population with generally low intake. To ensure population-level inference remains valid despite this skewed distribution, all analyses incorporated the IHS 2023 complex survey design and sampling weights, preserving national representativeness of estimated associations and standard errors.

Our findings underscore the necessity for health promotion strategies aimed at increasing fruit and vegetable consumption to be specifically tailored to distinct socio-demographic groups. Key factors warranting consideration for targeted interventions include gender, education level, employment status, economic status, marital status, place of residence, and area classification. Priority for interventions should be given to individuals who are male, possess lower educational attainment, are unemployed, belong to higher economic status groups, are unmarried, reside in regions such as Sumatra, Kalimantan, or Bali, and live in urban areas.

Gender disparities in dietary behavior are particularly salient, with women consistently demonstrating a higher propensity to consume fruits and vegetables compared to men. This observation aligns with previous studies reporting significantly higher intake levels among women across various contexts [7, 9–11]. For instance, Li et al [11] found that women in China consumed more fruit than men, and similar patterns were observed in Turkey, where women exhibited higher fruit and vegetable consumption than their male counterparts [7]. A consistent positive correlation was observed between higher educational attainment and healthier dietary

Tab. II. Results of the Analysis on the Association Between Sociodemographic Factors and Fruit and Vegetable Consumption in Indonesia.

Variables	Fruit and vegetable diet (%)		OR (CI 95%)	aOR (CI 95%)
	Less Healthy	Healthy		
Age				
19-59 years	83.6	83.1	Reference	
≥60 years	16.4	16.9	1.07 (1.00-1.14)	
Gender				
Male	45.4	41.6	Reference	Reference
Female	54.6	58.4	1.23 (1.18-1.30)*	1.31 (1.24-1.38)*
Education level				
No formal education	5.1	3.8	Reference	Reference
Primary education	50.5	41.9	1.03 (0.88-1.20)	0.99 (0.84-1.15)
Secondary education	32.4	34.5	1.36 (1.16-1.61)*	1.16 (0.98-1.37)
Higher education	12.0	19.8	2.29 (1.93-2.72)*	1.47 (1.23-1.75)*
Employment status				
Unemployed	33.1	31.9	Reference	Reference
Employed	66.9	68.1	1.06 (1.01-1.11)*	1.13 (1.06-1.20)*
Economic status				
Lowest	19.6	32.9	Reference	Reference
Lower-middle	20.2	20.9	0.57 (0.52-0.62)*	0.62 (0.56-0.68)*
Middle	19.9	16.4	0.45 (0.41-0.50)*	0.50 (0.46-0.56)*
Upper middle	20.1	15.4	0.44 (0.39-0.48)*	0.49 (0.44-0.54)*
Highest	20.2	14.4	0.42 (0.37-0.47)*	0.45 (0.40-0.51)*
Marital status				
Unmarried	21.2	17.5	Reference	Reference
Married	78.8	82.5	1.29 (1.20-1.39)*	1.25 (1.16-1.34)*
Place of residence				
Jawa	28.3	32.7	Reference	Reference
Sumatra	31.2	24.8	0.85 (0.76-0.95)*	0.85 (0.76-0.95)*
Kalimantan	9.5	7.1	0.82 (0.71-0.95)*	0.81 (0.70-0.94)*
Sulawesi	14.9	15.8	1.05 (0.93-1.18)	1.07 (0.95-1.22)
Bali	2.3	2.1	0.93 (0.76-1.14)	0.81 (0.66-0.99)*
Nusa Tenggara	6.0	7.6	1.18 (1.02-1.36)*	1.32 (1.13-1.53)*
Maluku	3.4	4.1	1.43 (1.10-1.90)*	1.54 (1.16-2.06)*
Papua	4.2	5.8	1.95 (1.46-2.60)*	1.95 (1.46-2.60)*
Area Classification				
Urban	53.4	59.1	Reference	Reference
Rural	46.6	40.9	0.86 (0.79-0.95)*	1.12 (1.02-1.23)*

* *p-value* < 0.05. Source: IHS 2023

patterns. This trend is consistent with other research indicating that education level significantly influences fruit and vegetable consumption [7, 12-14]. Consequently, efforts to enhance fruit and vegetable intake could be prioritized at the individual level, targeting those with no formal education, followed sequentially by individuals with primary, secondary, and higher educational backgrounds. Health professionals are further advised to develop health promotion strategies that are specifically tailored to the educational characteristics of their target groups.

Our study also found that unemployed individuals were less likely to consume fruits and vegetables, a result corroborated by previous research [13-15]. Employment status can affect fruit and vegetable intake, as individuals in formal employment often have stable incomes and better access to healthy foods, while unemployed

individuals may face financial constraints that limit their ability to purchase such foods [7].

Interestingly, the relationship between economic status and fruit and vegetable consumption in this study diverged from previous findings, which typically report higher intake among wealthier individuals [7, 10, 11]. Here, individuals in the highest economic group consumed fewer fruits and vegetables than those in the lowest economic group. This discrepancy may be explained by the measurement of economic status, which in this study was based on ownership of durable goods rather than actual disposable income. Thus, a materially affluent individual may not necessarily translate to a healthier diet, as intake is influenced by factors beyond purchasing power, such as time constraints, personal preferences, social environment, and lifestyle choices. This inverse association between higher economic

status (as measured by durable goods) and lower fruit/vegetable intake highlights a modern lifestyle trend that warrants further investigation to better understand consumer behavior.

Married individuals exhibited a greater tendency to consume fruits and vegetables compared to their unmarried counterparts, a finding consistent with previous research showing higher intake among married populations [7, 13, 16]. This association may be explained by shared meal routines, spousal encouragement, and household food decisions that support healthier eating patterns. Although these mechanisms were not directly assessed in the present study, they offer a reasonable explanation for the observed relationship.

Geographical disparities in fruit and vegetable consumption were also evident, with regions such as Sulawesi, Nusa Tenggara, Maluku, and Papua demonstrating higher proportions of healthy diets compared to Java. This suggests that less developed areas like Nusa Tenggara and Papua consume more fruits and vegetables than more developed regions such as Sumatra, Kalimantan, and Java [12]. Place of residence influences eating patterns through factors such as access to healthy food, prevailing community habits, and local food availability. These findings emphasize the critical need for context-specific interventions that are tailored to regional and population characteristics. Policymakers should leverage evidence-based data to formulate targeted strategies aimed at reducing dietary disparities across regions and social strata.

Within the framework of the Social Determinants of Health (SDH), sociodemographic factors – including gender, education, occupation, economic status, and geographic location – act as social determinants that can profoundly influence individual health behaviors, such as fruit and vegetable consumption. Discrepancies in sociodemographic characteristics, access to resources, health knowledge, and social environments collectively illustrate how these social determinants contribute to disparities in public health outcomes. Consequently, the findings of this study provide a foundational basis for designing multi-level, equity-oriented intervention strategies.

Given that inadequate fruit and vegetable consumption is a well-established risk factor for NCDs, dedicated attention is required to promote healthier diets across all segments of society. This aligns with recommendations from other studies, which identify the promotion of fruit and vegetable intake as a pivotal strategy in the prevention and control of NCDs.

To foster improved dietary patterns, particularly in terms of fruit and vegetable consumption, multi-level interventions are indispensable. These interventions should encompass educational initiatives, environmental modifications, and supportive public policies. These align with global evidence linking low fruits and vegetables intake to increased NCDs burden [17, 18], justifying targeted interventions for our identified sociodemographic disparities. Such efforts should be implemented gradually, coherently, and sustainably, as

recommended by Woodside et al. [19]. By integrating individual, community, and structural approaches, it is anticipated that the prevalence of low fruit and vegetable intake in Indonesia can be minimized, thereby contributing to a healthier and more equitable society.

The study identified a particularly vulnerable group with the lowest fruit and vegetable intake, comprising males, individuals without formal education, the unemployed, those in higher economic strata (as measured by durable goods), unmarried individuals, and residents of Sumatra, Kalimantan, Bali, and urban areas. Their heightened vulnerability stems from limited access, lower motivation, reduced awareness, specific social roles, and insufficient environmental support. Addressing these disparities necessitates robust multi-sectoral collaboration. For instance, the Ministry of Agriculture could facilitate the development of community gardens or affordable fruit stalls. The National Food Agency could provide fresh vegetable packages to low-income families. The Ministry of Manpower and the private sectors could integrate nutrition education into labor programs. The Ministry of Higher Education, Science, and Technology should engage local leaders as change agents in nutrition education. Health promotion campaigns can be bolstered through peer support and the involvement of male role models who actively practice healthy eating, perhaps through culturally sensitive campaigns that resonate with the target demographic.

The government plays a crucial role in supporting fruit and vegetable consumption through price regulations and market governance. To incentivize healthy eating, policies could include subsidizing local horticultural products, taxing highly processed foods, and regulating healthy food distribution in public institutions. These policy interventions are essential to counteract prevailing market trends that often favor inexpensive, calorie-dense, and nutrient-poor foods [20, 21]. Therefore, beyond educational initiatives, policy-driven structural strategies are paramount to ensuring equitable and sustainable health promotion outcomes.

The measurement of economic status using household durable goods ownership in the IHS dataset revealed a counterintuitive association: higher economic status correlated with lower fruit and vegetable consumption. This type of measurement, based on static assets, may not accurately reflect an individual's current, fluid purchasing ability or disposable income for daily dietary choices. This suggests that increased wealth, when assessed by durable goods, does not automatically translate to healthier diets, as intake is influenced by a complex interplay of factors beyond purchasing power, including time constraints, personal preferences, social environment, and modern lifestyle choices. This specific modern lifestyle trend merits further dedicated research to gain a deeper understanding of consumer behavior.

Rising malnutrition rates amplify global concerns of obesity and overweight as a pandemic. These findings suggest that interventions to improve fruit and vegetables consumption should be tailored to the sociodemographic groups identified in this study as having lower intake.

Since eating behavior is influenced by personal motivations, physiological preferences, and socio-cultural factors across all age groups, tailored efforts are needed to meet these diverse needs. For example, visual education and peer support can effectively engage unemployed or single men, while subsidies or community fruit markets may help those with lower incomes. Such focused approaches are crucial to closing consumption gaps among vulnerable populations [22-24].

Finally, observations from Maluku underscore that fruit and vegetable consumption is shaped by a confluence of individual, social, and spatial factors. In rural Maluku, many households possess fruit trees (*e.g.*, mangoes), providing direct and immediate access to fruits [25]. This contrasts with urban areas, where limited land availability reduces local food sources and increases reliance on market purchases. Our findings indicate higher consumption in rural areas, while urban areas showed no significant association. This highlights the imperative to consider spatial and ecological factors in nutrition policies, advocating for initiatives such as supporting urban farming, promoting home gardens, and revitalizing traditional markets to boost fruit and vegetable intake in urban settings.

Based on the observed associations, future interventions should be tailored to the sociodemographic groups identified in the present study, with emphasis on improving access to and awareness of fruit and vegetables consumption among populations at higher risk of inadequate intake.

Conclusion

Overall, 97.1% of Indonesian adults did not meet the WHO-STEPS fruit and vegetable intake threshold (≥ 5 servings/day for 7 consecutive days). Although this stringent benchmark produces a highly imbalanced classification in a generally low-intake setting, it ensures international comparability and helps identify priority groups for targeted dietary interventions. Promotion and intervention strategies to increase fruit and vegetable consumption should be tailored and designed based on the priority target groups, specifically those with low intake or considered vulnerable. Vulnerable groups include males, individuals aged 19-59 years, those with low education, the unemployed, and unmarried individuals. For example, initiatives may involve encouraging school canteen vendors to provide fruits and vegetables, inviting fruit sellers to offer pre-cut fruit at schools, and conducting outreach programs on the importance of fruit consumption, particularly targeting unmarried men. In addition, advocacy is needed to gain policy support from relevant stakeholders.

Acknowledgements

None.

Funding

None.

Ethical Approval

This study received ethical approval from the Medical and Health Research Ethics Committee of the Faculty of Medicine, Public Health, and Nursing, Universitas Gadjah Mada, under permit number KE/FK/0117/EC/2025.

Conflict of interest statement

The authors have no conflicts of interest to report.

Authors' contributions

AA contributed to the study conceptualization, data curation, formal analysis, interpretation of findings, and drafting of the manuscript as part of her postgraduate study. ALR and FMK supervised the study, contributed to the study design and methodology, provided academic guidance throughout the research process, and critically reviewed and revised the manuscript. DNP and AW served as examiners, contributed to methodological and analytical feedback, and critically reviewed the manuscript. All authors read and approved the final version of the manuscript.

References

- [1] Noncommunicable Diseases (NCD) Country Profiles 2018. Geneva: World Health Organization; 2018. Licence: CC BY-NC-SA 3.0 IGO. Available at: <https://www.who.int/publications/item/9789241514620> Accessed on: 29/10/2024.
- [2] Directorate General of Disease Prevention and Control, Ministry of Health, Republic of Indonesia. 2023. Annual Report 2022 Prevention and Control of Non-Communicable Diseases (NCDs).
- [3] Purnamasari D. The Emergence of Non-communicable Disease in Indonesia. *Acta Med Indones* 2018;50(4):273-4. Available at: <https://actamedindones.org/index.php/ijim/article/view/1028>. Accessed on: 24 July 2024.
- [4] Neuwald N V., Pearce AL, Adise S, Rolls BJ, Keller KL. Switching between foods: A potential behavioral phenotype of hedonic hunger and increased obesity risk in children. *Physiol Behav* 2023;270:114312. <https://doi.org/10.1016/j.physbeh.2023.114312>.
- [5] Burke DT, Bennett AE, Hynds P, Priyadarshini A. Identifying Novel Data-Driven Dietary Patterns via Dimensionality Reduction and Associations with Socioeconomic Profile and Health Outcomes in Ireland. *Nutrients* 2023;15:3256. <https://doi.org/10.3390/nu15143256>.
- [6] Jideani AIO, Silungwe H, Takalani T, Omolola AO, Udeh HO, Anyasi TA. Antioxidant-rich natural fruit and vegetable products and human health. *Int J Food Prop* 2021;24:41-67. <https://doi.org/10.1080/10942912.2020.1866597>.
- [7] Küçük N, Urak F, Bilgic A, Florkowski WJ, Kiani AK, Özdemir FN. Fruit and vegetable consumption across population segments: evidence from a national household survey. *J Heal*

- Popul Nutr 2023;42:1-20. <https://doi.org/10.1186/s41043-023-00382-6>.
- [8] Badan Kebijakan dan Pembangunan Kesehatan. Survei Kesehatan Indonesia 2023 dalam Angka. Jakarta: Kementerian Kesehatan Republik Indonesia 2023.
 - [9] Lee-Kwan SH, Moore LV, Blanck HM, Harris DM, Galuska D. Disparities in State-Specific Adult Fruit and Vegetable Consumption - United States, 2015. *MMWR Morb Mortal Wkly Rep* 2017;66:1241-7. <https://doi.org/10.15585/mmwr.mm6645a1>.
 - [10] Hoy MK, Goldman JD, Moshfegh AJ. Differences in fruit and vegetable intake of U.S. adults by sociodemographic characteristics evaluated by two methods. *J Food Compos Anal* 2017;64:97-103. <https://doi.org/10.1016/j.jfca.2017.06.012>.
 - [11] Li L, Ouyang Y, Wang H, Huang F, Wang Y, Zhang J, Su C, Du W, Jia X, Jiang H, Wang Z, Zhang B. Disparities in fresh fruit and vegetable intake by sociodemographic and behavioural factors among adults in China. *Public Health Nutr* 2022;25:649-56. <https://doi.org/10.1017/S1368980020003274>.
 - [12] Darmawan ES, Kusuma D, Permanasari VY, Amir V, Tjandrarini DH, Dharmayanti I. Beyond the Plate: Uncovering Inequalities in Fruit and Vegetable Intake across Indonesian Districts. *Nutrients* 2023;15:1-13. <https://www.mdpi.com/2072-6643/15/9/2160>.
 - [13] Affret A, Severi G, Dow C, Mancini FR, Rey G, Delpierre C, Clavel-Chapelon F, Boutron-Ruault M, Fagherazzi G. Socio-economic factors associated with an increase in fruit and vegetable consumption: A 12-year study in women from the E3N-EPIC study. *Public Health Nutr* 2017;21:740-55. <https://doi.org/10.1017/S1368980017003196>.
 - [14] Craveiro D, Marques S, Zverinova I, Maca V, Milan S, Chiabai A, Suarez C, Martinez-Juarez P, Jalon SG, Quiroga S, Taylor T. Explaining inequalities in fruit and vegetable intake in Europe: The role of capabilities, opportunities and motivations. *Appetite* 2021;165:105283. <https://doi.org/10.1016/j.appet.2021.105283>.
 - [15] Giacomuzzo B, Ngo MH, Ha TM, Markoni E, Delley M, Götze F, Le NT, Bui TL, Nguyen AD, Pham BD, Brunner TA, Lucas BF. Determinants of the intention to increase vegetable consumption in Vietnam and Switzerland. *J Agric Food Res* 2024;15:4-8. <https://doi.org/10.1016/j.jafr.2024.100998>.
 - [16] Vinther JL, Conklin AI, Wareham NJ, Monsivais P. Marital transitions and associated changes in fruit and vegetable intake: Findings from the population-based prospective EPIC-Norfolk cohort, UK. *Soc Sci Med* 2016;157:120-6. <http://dx.doi.org/10.1016/j.socscimed.2016.04.004>.
 - [17] Smith L, López Sánchez GF, Veronese N, Soysal P, Oh H, Barnett Y, Keyes H, Butler L, Allen P, Kostev K, Jacob L, Shin J.I, Koyanagi A. Fruit and Vegetable Intake and Non-Communicable Diseases among Adults Aged ≥50 Years in Low- and Middle-Income Countries. *J Nutr Heal Aging* 2022;26:1003-9. <https://doi.org/10.1007/s12603-022-1855-z>.
 - [18] Afshin A, Sur PJ, Fay KA, Cornaby L, Ferrara G, Salama JS, Mul-lany EC, Abate KH, Abbafati C, Abebe Z, Afarideh M, Aggarwal A, Agrawal S, Akinyemiju T, Alahdab F, Bacha U, Bachman VF, Badali H, Badawi A, Bensenor IM, Bernabe E, Biryukov SH, Biadgilign SKK, Cahill LE, Carrero JJ, Cercy KM, Dandona L, Dandona R, Dang AK, Degefa MG, El Sayed Zaki M, Esteghamati A, Esteghamati S, Fanzo J, Farinha CSE, Farvid MS, Farzad-far F, Feigin VL, Fernandes JC, Flor LS, Foigt NA, Forouzanfar MH, Ganji M, Geleijnse JM, Gillum RF, Goulart AC, Grosso G, Guessous I, Hamidi S, Hankey GJ, Harikrishnan S, Hassen HY, Hay SI, Hoang CL, Horino M, Ikeda N, Islami F, Jackson MD, James SL, Johansson L, Jonas JB, Kasaeian A, Khader YS, Khalil IA, Khang YH, Kimokoti RW, Kokubo Y, Kumar GA, Lallukka T, Lopez AD, Lorkowski S, Lotufo PA, Lozano R, Malekzadeh R, März W, Meier T, Melaku YA, Mendoza W, Mensink GBM, Michal R, Miller TR, Mirarefin M, Mohan V, Mokdad AH, Mozaffarian D, Nagel G, Naghavi M, Nguyen CT, Nixon MR, Ong KL, Pereira DM, Poustchi H, Qorbani M, Rai RK, Razo-García C, Rehm CD, Rivera JA, Rodríguez-Ramírez S, Roshandel G, Roth GA, Sanabria J, Sánchez-Pimienta TG, Sartorius B, Schmidhuber J, Schutte AE, Sepanlou SG, Shin MJ, Sorensen RJD, Springmann M, Szponar L, Thorne-Lyman AL, Thrift AG, Touvier M, Tran BX, Tyrovolas S, Ukwaja KN, Ullah I, Uthman OA, Vaezghasemi M, Vasankari TJ, Vollset SE, Vos T, Vu GT, Vu LG, Weiderpass E, Werdecker A, Wijeratne T, Willett WC, Wu JH, Xu G, Yone-moto N, Yu C, Murray CJL. Health effects of dietary risks in 195 countries, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 2019;393:1958-72. [http://dx.doi.org/10.1016/S0140-6736\(19\)30041-8](http://dx.doi.org/10.1016/S0140-6736(19)30041-8).
 - [19] Woodside J V., Nugent AP, Moore RE, McKinley MC. Fruit and vegetable consumption as a preventative strategy for non-communicable diseases. *Proc Nutr Soc* 2023;186-99. <https://doi.org/10.1017/S0029665123002161>.
 - [20] Reeve E, Ravuvu A, Johnson E, Nasiga S, Brewer T, Mounsey S, Thow AM. Scaling up food pricing policies in the Pacific: A guide to action. *BMJ Glob Heal* 2023;8:8-10. <http://dx.doi.org/10.1136/bmjgh-2023-012041>.
 - [21] WHO. Fiscal policies to promote healthy diets: WHO guideline. 2024, pp. 1-24. Available at: <https://www.who.int/publications/i/item/9789240091016>. Accessed on: 21/08/2024.
 - [22] Agurs-Collins T, Alvidrez J, ElShourbagy Ferreira S, Evans M, Gibbs K, Kowtha B, Pratt C, Reedy J, Shams-White M, Brown AGM. Perspective: Nutrition Health Disparities Framework: A Model to Advance Health Equity. *Adv Nutr* 2024;15:100194. <https://doi.org/10.1016/j.advnut.2024.100194>.
 - [23] Ciangherotti S, Kevicka, K, Ciangherotti E, Rondinone R. Eating behaviour: Research in Liguria on young, adult and elderly subjects. *J Prev Med Hyg* 2006;47:26-36. <https://doi.org/10.15167/2421-4248/jpmh2006.47.1.13>.
 - [24] Crovetto M, Sepúlveda MJ. Relationship between dietary energy intake, nutritional status and cardiovascular risk in adults from the communes of Quellón and Chonchi, Chiloé, Chile. *J Prev Med Hyg* 2022;63:E435-E441. <https://doi.org/10.15167/2421-4248/jpmh2022.63.3.2486>.
 - [25] Sitepu AE, Pakpahan M, Cicilia SL. The Determinant Impacting Fruit and Vegetable Consumption Among Families in Maluku, Indonesia. *Indones J Public Heal* 2025;20:70-81. <https://doi.org/10.20473/Ijph.v20i1.2025.70-81>.

Received on September 24, 2026. Accepted on April 30, 2026.

Correspondence: Aditya Lia Ramadana, Department of Health Behaviour, Environment, and Social Medicine, Faculty of Medicine, Public Health, and Nursing, Universitas Gadjah Mada, Gedung S2 IKM Lantai 3, Jl. Farmako Sekip Utara Yogyakarta 55281, Indonesia. E-mail: al-ramadana@ugm.ac.id.

How to cite this article: Amini A, Kusumaningrum FM, Purwaningrum DN, Wahab A, Ramadana AL. Sociodemographic determinants of fruit and vegetable consumption in Indonesia: insights from the 2023 Indonesian Health Survey. *J Prev Med Hyg* 2026;67:E188-E194. <https://doi.org/10.82082/2421-4248/jpmh2026.67.2.3749>

© Copyright by Pacini Editore Srl, Pisa, Italy

This is an open access article distributed in accordance with the CC-BY-NC-ND (Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International) license. The article can be used by giving appropriate credit and mentioning the license, but only for non-commercial purposes and only in the original version. For further information: <https://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>