

ORIGINAL ARTICLE

Emergence of plasmid-mediated quinolone-resistant determinants in *Klebsiella pneumoniae* isolates from Tehran and Qazvin provinces, Iran

A. PEYMANI¹, T. NASERPOUR FARIVAR¹, L. NIKOOEI¹, R. NAJAFIPOUR¹, A. JAVADI², A.A. PAHLEVAN¹¹ Cellular and Molecular Research Center, Qazvin University of Medical Sciences, Qazvin, Iran; ² School of Allied Sciences, Tehran University of Medical Sciences, Tehran, Iran

Key words

Klebsiella pneumoniae • Quinolones resistance • *qnr*

Summary

Background. Plasmid-mediated quinolone resistance is an increasing clinical concern, globally. The major objective of the present study was to identify the *qnr*-encoding genes among the quinolone non-susceptible *K. pneumoniae* isolates obtained from two provinces in Iran.

Methods. A total of 200 *K. pneumoniae* isolates were obtained from hospitals of Qazvin and Tehran, Iran. The identification of bacterial isolates was carried out by standard laboratory methods and API 20E strips. Susceptibility to quinolone compounds were examined by standard Kirby-Bauer disk diffusion method according to the CLSI guideline. PCR and sequencing were employed to detect *qnrA*, *qnrB* and *qnrS*-encoding genes.

Results. Of 200 *K. pneumoniae* isolates, 124 (62%) were non-susceptible to quinolone compounds among those 66 (53.2%) and 58 (46.8%) isolates showed high and low-level quinolone resistance rates, respectively. Out of 124 quinolone non-susceptible isolates, *qnr*-encoding genes were present in 49 (39.5%) isolates with *qnrB1* (30.6%) as the most dominant gene followed by *qnrB4* (9.7%), and *qnrS1* (1.6%) either alone or in combination.

Conclusions. This study, for the first time, revealed the high appearance of *qnrB1*, *qnrS1* and *qnrB4* genes among the clinical isolates of *K. pneumoniae* in Iran. Therefore, the application of proper infection control measures and well-established antibiotic administration guideline should be strictly considered within our medical centers.

Introduction

Klebsiella pneumoniae (*K. pneumoniae*) is an opportunistic pathogen causing several nosocomial infections such as urinary tract infections, pneumonia, septicemia, and soft tissue infections [1]. This organism is also known as a community-acquired potential pathogen [2]. Health care associated infection caused by this organism has been linked to high mortality and morbidity especially among the patients admitted to intensive care units [3, 4].

Quinolones are a group of synthetic antibacterial agents that are widely used in routine clinical practice [5]. The new quinolones compounds (6-fluoroquinolones) exhibit broad spectrum of antibacterial activity against Gram-negative, mycobacterial pathogens, and anaerobes. Moreover, these agents show a good-to-moderate oral absorption and tissue penetration with favorable pharmacokinetics in humans, creating desirable clinical efficacy in treating many kinds of infections [6, 7]. Quinolones inhibit the function of bacterial DNA gyrase and topoisomerase IV [8]. While the first and second generation fluoroquinolones selectively inhibit the topoisomerase II ligase domain or DNA gyrase activity, the quinolones of third and fourth generations are with more tendency for topoisomerase IV ligase [9]. Excessive and

inappropriate administration of antimicrobial agents such as quinolones has increased the emergence of multidrug resistant *K. pneumoniae* isolates which makes the process of antimicrobial therapy to become marginal and problematic [10, 11]. In recent years, several studies have demonstrated that the appearance of quinolone-resistant *K. pneumoniae* is rising at a faster rate, worldwide [12-15]. Infections caused by resistant organisms are often due to extensive cross-resistance with other antimicrobials, including beta-lactams and aminoglycosides [16]. Quinolone resistance in *Enterobacteriaceae* mainly occurs through chromosomal mutations in the genes coding for DNA gyrase and topoisomerase IV, changes in outer membrane and efflux proteins or in their regulatory mechanisms [17]. Findings from recent studies show that plasmid-mediated resistance, associated with the pentapeptide proteins of the *qnr* family, might play a crucial role in quinolone compound resistance [18]. Three major groups of *qnr* determinants, *qnrA*, *qnrB*, and *qnrS*, are increasingly being identified in the clinical isolates of various enterobacterial species, worldwide [19]. It was in 1998 that the first plasmid-mediated quinolone resistance determinant, *qnrA*, was reported in a *Klebsiella pneumoniae* strain from the United States [20]. Since then two *qnr* determinants, *qnrB* and *qnrS* have been discovered in other *Enterobacte-*

riaceae species such as *Citrobacter Koseri*, *Escherichia coli*, *Enterobacter cloacae*, and *Klebsiella pneumoniae* from Asia and Europe [21-24]. To date, there has been no report for the frequency of *qnr* genes among *K. pneumoniae* isolates in Iran. In the current study, for the first time, we described the frequency of *qnr* determinants (*qnrA*, *qnrB*, and *qnrS*) among the isolates of quinolone non-susceptible *K. pneumoniae* collected from hospitals of Qazvin and Tehran provinces.

Methods

BACTERIAL ISOLATES AND ANTIMICROBIAL SUSCEPTIBILITY

In this descriptive study, a total of 200 clinical isolates of *K. pneumoniae* were collected from hospitalized patients in several teaching hospitals in Tehran and Qazvin during 2012-2013. The isolates were obtained from different clinical specimens including urine, wound, trachea, secretions, blood, and ascites. All isolates were identified by standard laboratory methods and confirmed with the API 20 E (bioMérieux, France) strips. All isolates were kept at -70°C in trypticase soy broth containing 20% glycerol and subcultured twice before testing. The mean age of patients (77 (38.5%) male and 123 (61.5%) female) was 51.7±17.4 (range 17-83) years. Written informed consent was obtained from all subjects enrolled in this study. Kirby-Bauer disk diffusion technique was performed according to the CLSI guideline to identify quinolone resistance using nalidixic acid (30 µg), ciprofloxacin (5 µg), gatifloxacin (5 µg), norfloxacin (10 µg), and levofloxacin (5 µg) disks [25]. In this study the isolates were classified either as high-level quinolone resistant if the resistance to both nalidixic acid and ciprofloxacin disks was observed or low-level quinolone resistant in the cases of resistance to nalidixic acid, presence of intermediate isolates or ciprofloxacin-susceptible organisms [26]. Antibiotic disks were purchased from the Mast (Mast Diagnostics Group Ltd, Merseyside, UK). *E. coli* ATCC 25922 and *Pseudomonas aeruginosa* ATCC 27853 were used as quality control strains in antimicrobial susceptibility testing.

DETECTION OF *QNR* DETERMINANTS

Detection of *qnrA*, *qnrB*, and *qnrS* plasmid-mediated quinolone resistance genes was performed using PCR

and specific primers (Tab. I). Plasmid DNA was extracted by plasmid mini extraction kit (Bioneer Company, South Korea). PCR amplifications were applied in a thermocycler (Applied Biosystems, USA) as follows: 95°C for 5min and 35 cycles of 1min at 95°C, 1min at specific annealing temperature for each primer and 1min at 72°C. A final extension step of 10 min at 72°C was performed. Amplification reactions were prepared in a total volume of 25 µl (24 µl of PCR master mix plus 1 µl of template DNA) including 5ng of genomic DNA, 2.0U of Taq DNA polymerase, 10mM dNTP mix at a final concentration of 0.2mM, 50mM MgCl₂ at a final concentration of 1.5mM, 1 µM of each primer, and 1X PCR buffer (final concentration). PCR products were electrophoresed on 1% agarose gel at 100 volts and later stained with ethidium bromide solution and finally visualized in a gel documentation system (UVtec, UK). The purified PCR products were sequenced by the Macrogen Company (Seoul, South Korea) and the sequence alignment and analysis were performed online using the BLAST program of the National Center for Biotechnology Information (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>).

Data were summarized using mean ± SD (standard deviation), proportional frequency and confidence interval for microbiological, clinical, and demographic characteristics. All analyses were carried out using a Statistical Software Package, SPSS for windows version 16.0 (Chicago, IL, USA).

Results

In this study, the bacterial isolates were recovered from different clinical specimens including urine (110-55.0%), trachea (59-29.5%), wound (18-9.0%), blood (8-4%), and ascites (5-2.5%). These isolates were obtained from the patients admitted to intensive care units (96-48.0%), internal medicine (54-27.0%), infectious diseases (35-17.5%), surgery (13-6.5%), and orthopaedic (2-1.0%) wards. The results of antimicrobial susceptibility testing showed the resistance rates against the antimicrobial agents used in our study varied between 20% and 58%. Overall, nalidixic acid (58%) and ciprofloxacin (34.5%) revealed the highest rates of resistance among the antimicrobials tested whereas levofloxacin and norfloxacin also demonstrated high susceptibility rates of 80% and 77%, respectively (Tab. II). In total,

Tab. I. The primers used for detection of *qnr* genes in *K. pneumoniae* isolated from Qazvin and Tehran hospitals.

PCR targets	Primer sequence (5'-3')	Annealing temperatures (°C)	References
qnrA1-6	F: ACGCCAGGATTTGAGTGAC R: CCAGGCACAGATCTTGAC	49	27
qnrB1-3, 5, 6, 8	F: GGCACCTGAATTATCGGC R: TCCGAATTGGTCAGATCG	49	27
qnrB4	F: AGTTGTGATCTCTCCATGGC R: CGGATATCTAAATCGCCAG	53	27
qnrS1-2	F: CCTACAATCATACATATCGGC R: GCTTCGAGAATCAGTCTTGC	53	27

Tab. II. Antibiotic susceptibility of *K. pneumoniae* against quinolone compounds.

Antimicrobial agents	Resistance n (%) [CI]	Intermediate n (%) [CI]	Susceptible n (%) [CI]
Nalidixic acid	83(41.5) [34.7-48.3]	33(16.5) [11.4-21.6]	84(42) [35.2-48.8]
Ciprofloxacin	53(27) [20.8-33.2]	15(7.5) [3.8-11.2]	131(65.5) [58.9-72.1]
Gatifloxacin	39(19.5) [14-25]	27(13.5) [8.8-18.2]	134(67) [60.5-73.5]
Norfloxacin	37(18.5) [13.1-23.9]	9(4.5) [1.6-7.4]	154(77) [71.2-82.8]
Levofloxacin	36(18) [12.7-23.3]	4(2) [0.1-3.9]	160(80) [74.5-85.5]

CI = 95% Confidence interval

66 (53.2%) and 58 (46.8%) of isolates showed high and low-level quinolone resistance, respectively.

PCR and sequencing showed the presence of *qnr*-encoding genes in 49 (39.5%) of quinolone non-susceptible *K. pneumoniae* isolates among those *qnrB1* (38-30.6%) was the most common gene followed by *qnrB4* (12-9.7%) and *qnrS1* (2-1.6%) genes either alone or in combination. The study isolates were negative for *qnrA* gene. As shown in Table III, *qnrB1* was found to coexist with *qnrB4* in 3 (2.4%) isolates. Overall, 25 (37.9%) high level quinolone resistant isolates carried *qnr* genes in which 19 (28.8%), 4 (6.1%), and 2 (3%) isolates carried *qnrB1*, *qnrB4*, and *qnrS1* genes, respectively. In ad-

Tab. III. Distribution of *qnrB1*, *qnrB4*, and *qnrS1* genes among *qnr*-positive *K. pneumoniae* isolates.

<i>qnr</i> -encoding genes	N of isolates n (%) [CI]
<i>qnrB1</i>	35 (28.2%) [20.3-36.1]
<i>qnrB4</i>	9 (7.3%) [2.7-11.9]
<i>qnrS1</i>	2 (1.6%) [0-3.8]
<i>qnrB1+qnrB4</i>	3 (2.4%) [0-5.1]
<i>qnr</i> negative	75(60.5%) [51.9-69.1]
Total	124 (100%)

CI: 95% Confidence interval

Tab. IV. Frequency of *qnr*-positive *K. pneumoniae* isolates based on hospital wards and source of Specimens (n = 49).

Wards	N° of isolates n (%) [CI]	Specimens	N° of isolates n (%) [CI]
ICU	29 (59.2%) [45.4-73]	Urine	21 (42.9%) [29-56.8]
Internal medicine	10 (20.4%) [9.1-31.7]	Trachea	18 (36.7%) [23.2-50.2]
Infectious diseases	8 (16.3%) [6-26.6]	Wound	6 (12.2%) [3-21.4]
Surgery	2 (4.1%) [0-9.7]	Blood	1 (2%) [0-5.9]
Orthopedic	-	Ascites	3 (6.1%) [0-12.8]

ICU: Intensive Care Unit

CI: 95% Confidence interval

dition, 24 (41.4%) low level quinolone resistant isolates were positive for *qnr* genes among those 19 (32.8%) isolates carried *qnrB1* gene followed by *qnrB4* in 8 (13.8%) isolates. Among the high and low-level quinolone resistance isolates, *qnrB1* was the most frequent gene compared to other genes. Table IV shows that *qnr*-positive isolates were mostly recovered from urine (42.9%) followed by trachea secretion (36.7%) samples. The patients affected by these organisms were mostly admitted to ICU (59.2%) and internal medicine (20.4%) wards.

Discussion

K. pneumoniae is being increasingly recognized as a clinically significant nosocomial pathogen [1]. Quinolones are among the most commonly administered antimicrobials routinely used for the treatment of serious infections caused by *K. pneumoniae* and other members of the genus Enterobacteriaceae [6]. However, the development of resistance to these antibiotics makes the treatment decision difficult, leading to treatment failures [5]. In recent years, plasmid mediated quinolone resistance among enterobacterial isolates has been reported in several studies, worldwide. However, the number of reports on prevalence of *qnr* genes among Iranian enterobacteria isolates is only limited to few studies [28, 29].

In the present study, 58% and 34.5% of isolates were fully or intermediate resistant to nalidixic acid and ciprofloxacin, respectively. These findings were higher than the two previously conducted studies in Iran. Raei et al demonstrated that 36.2% and 34.1% of urinary *K. pneumoniae* isolates were resistant to ciprofloxacin and nalidixic acid, respectively [30]. In another study from Iran, Zamani et al found that 28.57% and 23.8% of *Klebsiella* spp. were resistant to nalidixic acid and ciprofloxacin, respectively [31]. Hence, the emergence of resistant isolates against broad spectrum antibacterial agents in our hospital settings seems to be linked with improper and widespread administration of these antibiotics.

The present study demonstrates a high prevalence (39.5%) for plasmid-mediated quinolone resistance determinants among quinolone non-susceptible *K. pneumoniae* isolates in Iran. The prevalence rate found in our study is higher than those reported by Kim et al from Korea (10%) [32], Wang et al from China (11.9%) [33], Dahmen et al from Tunisia (16%) [34], Yan et al from China (16.2%) [35], and Wang et al from the United States (11.1%) [36] but still lower than that found by Bouchakour et al in Morocco in which 50% of ESBL-producing *K. pneumoniae* isolates were shown to carry *qnr* determinants [37]. This might be indicative of a rising trend in the rate of plasmid mediated quinolone resistance among the genus of Enterobacteriaceae.

In the current study, 25% of *qnr*-positive isolates were shown to have high level quinolone resistance. As plasmid mediated quinolone resistance determinants produce only low-level resistance to quinolones, it can be hypothesized that high level resistant pattern is possibly

caused by another mechanisms such as chromosomal mutation which was not evaluated in the present study. Considering the findings of the present study, it is obvious that most *qnr*-positive *K. pneumoniae* isolates were mostly obtained from the patients admitted to ICUs. Long term ICU stay, broad spectrum antibiotics intake, chronic underlying conditions, and the application of invasive techniques and devices probably make the patients more susceptible to infections caused by these resistant organisms.

In the present study, 30.6%, 9.7%, and 1.6% of quinolone non-susceptible *K. pneumoniae* isolates carried *qnrB1*, *qnrB4*, and *qnrS1* genes alone or in combination, respectively. We believe that this is the first report of *qnrS1*, *qnrB4*, and *qnrB1* genes among the clinical isolates of *K. pneumoniae* collected from two distinct provinces of Iran. In a study by Pakzad et al reported from Iran, 9 (37.5%) and 4 (20.8%) of ESBL-producing *E. coli* isolates were positive for *qnrA* and *qnrB* genes, respectively [29]. The presence of *qnrA* (25.8%), *qnrB1* (1.17%), and *qnrS* (1.17%) genes among ESBL-producing *Salmonella* spp. was also reported in a study by Saboohi et al from Iran [28]. In another study from Iran, Seyedpour et al showed that 30.4% of community isolates of *K. pneumoniae* harbored *qnr* and/or *aac* (6')-Ib-cr genes [38]. In Taiwan, Wu et al described the presence of *qnrB4* (3.6%), *qnrS1* (2.8), and *qnrB2* (2.3%) genes in the clinical isolates of *K. pneumoniae* [39]. Robicsek et al in the United States reported that 14% and 6% of ceftazidime-resistant *K. pneumoniae* isolates harbored *qnrA* and *qnrB* genes, respectively [40]. Dahmen et al from Tunisia showed *qnrA* was more prevalent among *K. pneumoniae* isolates whereas *qnrB1* was the most prevalent genes among *E. cloacae* isolates followed by *qnrB2* and *qnrS1* [34]. Similarly, Yan et al in their report from China demonstrated that 8.1%, 4.1%, and 4.1% of ESBL-producing *K. pneumoniae* isolates were positive for *qnrA*, *qnrB*, and *qnrS* genes, respectively [35]. Finally, Wang et al in a study carried out in China reported that 62 (15.1%), 25 (6.1%), and 10 (2.4%) of ESBL-producing *K. pneumoniae* isolates were positive for *qnrS*, *qnrB*, and *qnrA* genes, respectively [33].

Conclusions

Findings of the present study reveal a high prevalence for plasmid-mediated quinolones resistance due to *qnr* genes among the clinical isolates of *K. pneumoniae* in Iran. The appearance and spread of such resilient organisms within the medical centers around the country not only brings about issues of great concern for human health but also raises questions on how to achieve a successful antibiotic therapy through planning a comprehensive infection control guideline to avoid further spread of these resistant organisms within our medical settings. Our data also highlights the necessity for establishing an appropriate infection control strategy and sensible antibiotic therapy.

ACKNOWLEDGMENTS

This study was financially supported by the Cellular and Molecular Research Center and the Research Deputy of Qazvin University of Medical Sciences (grant number 20/5233), Qazvin, Iran.

References

- [1] Podschun R, Ullmann U. *Klebsiella* spp. as nosocomial pathogens: epidemiology, taxonomy, typing methods, and pathogenicity factors. Clin Microbiol Rev 1998;11:589-603.
- [2] Ko WC, Paterson DL, Sagnimeni AJ, et al. Community-acquired *Klebsiella pneumoniae* bacteremia: global differences in clinical patterns. Emerg Infect Dis 2002;8:160-6.
- [3] Orsi GB, d'Ettorre G, Panero A, et al. Hospital-acquired infection surveillance in a neonatal intensive care unit. Am J Infect Control 2009;37:201-3.
- [4] Bennett JW, Robertson JL, Hospenthal DR, et al. Impact of extended spectrum beta-lactamase producing *Klebsiella pneumoniae* infections in severely burned patients. J Am Coll Surg 2010; 211: 391-9.
- [5] Fàbrega A, Madurga S, Giralt E, et al. Mechanism of action of and resistance to quinolones. Microb Biotechnol 2009;2:40-61.
- [6] Emami S, Shafiee A, Foroumadi A. Quinolones: recent structural and clinical developments. Iran J Pharm Res 2010;4:123-36.
- [7] Wiles JA, Bradbury BJ, Pucci MJ. New quinolone antibiotics: a survey of the literature from 2005 to 2010. Expert Opin Ther Pat 2010;20:1295-319.
- [8] Drlica K, Malik M, Kerns RJ, et al. Quinolone-mediated bacterial death. Antimicrob Agents Chemother 2008;52:385-92.
- [9] Adams DE, Shekhtman EM, Zechiedrich EL, et al. The role of topoisomerase IV in partitioning bacterial replicons and the structure of catenated intermediates in DNA replication. Cell 1992;71:277-88.
- [10] Lautenbach E, Strom BL, Bilker WB, et al. Epidemiological investigation of fluoroquinolone resistance in infections due to extended-spectrum beta-lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae*. Clin Infect Dis 2001;33:1288-94.
- [11] Paterson DL, Mulazimoglu L, Casellas JM, et al. Epidemiology of ciprofloxacin resistance and its relationship to extended-spectrum beta-lactamase production in *Klebsiella pneumoniae* isolates causing bacteremia. Clin Infect Dis 2000;30:473-8.
- [12] Wang M, Sahm DF, Jacoby GA, et al. Emerging plasmid-mediated quinolone resistance associated with the *qnr* gene in *Klebsiella pneumoniae* clinical isolates in the United States. Antimicrob Agents Chemother 2004;48:1295-99.
- [13] Wang A, Yang Y, Lu Q, et al. Presence of *qnr* gene in *Escherichia coli* and *Klebsiella pneumoniae* resistant to ciprofloxacin isolated from pediatric patients in China. BMC Infect Dis 2008;8:68.
- [14] Yu WL, Jones RN, Hollis RJ, et al. Molecular epidemiology of extended-spectrum beta-lactamase-producing, fluoroquinolone-resistant isolates of *Klebsiella pneumoniae* in Taiwan. J Clin Microbiol 2002;40:466-9.
- [15] Rodríguez-Martínez JM, Pascual A, García I, et al. Detection of the plasmid-mediated quinolone resistance determinant *qnr* among clinical isolates of *Klebsiella pneumoniae* producing AmpC-type beta-lactamase. J Antimicrob Chemother 2003;52:703-6.
- [16] Tolun V, Küçükbasmacı O, Törümküney-Akbulut D, et al. Relationship between ciprofloxacin resistance and extended-spectrum beta-lactamase production in *Escherichia coli* and *Klebsiella pneumoniae* strains. Clin Microbiol Infect 2004;10:72-5.
- [17] Ruiz J. Mechanisms of resistance to quinolones: target alterations, decreased accumulation and DNA gyrase protection. J Antimicrob Chemother 2003;51:1109-17.

- [18] Robicsek A, Jacoby GA, Hooper DC. *The worldwide emergence of plasmid-mediated quinolone resistance*. Lancet Infect Dis 2006;6:629-40.
- [19] Cattoir V, Poirel L, Nordmann P. *Plasmid-mediated quinolone resistance determinant QnrB4 identified in France in an Enterobacter cloacae clinical isolate coexpressing a QnrS1 determinant*. Antimicrob Agents Chemother 2007;51:2652-3.
- [20] Martínez-Martínez L, Pascual A, Jacoby GA. *Quinolone resistance from a transferable plasmid*. Lancet 1998;351:797-9.
- [21] Jonas D, Biehler K, Hartung D, et al. *Plasmid-mediated quinolone resistance in isolates obtained in German intensive care units*. Antimicrob Agents Chemother 2005;49:773-5.
- [22] Mammeri H, Van De Loo M, Poirel L, et al. *Emergence of plasmid-mediated quinolone resistance in Escherichia coli in Europe*. Antimicrob Agents Chemother 2005;49:71-6.
- [23] Nazic H, Poirel L, Nordmann P. *Further identification of plasmid-mediated quinolone resistance determinant in Enterobacteriaceae in Turkey*. Antimicrob Agents Chemother 2005;49:2146-7.
- [24] Firoozeh F, Zibaei M, Soleimani-Asl Y. *Detection of plasmid-mediated qnr genes among the quinolone-resistant Escherichia coli isolates in Iran*. J Infect Dev Ctries 2014;8: 818-22.
- [25] Clinical and Laboratory Standards Institute. *Performance standards for antimicrobial susceptibility testing: seventeenth informational supplement M100-S17*. 2013; Wayne, PA, USA.
- [26] Oktem IM, Gulay Z, Bicmen M, et al. *qnrA prevalence in extended-spectrum beta-lactamase-positive Enterobacteriaceae isolates from Turkey*. Jpn J Infect Dis 2008;6:13-7.
- [27] Lavilla S, González-López JJ, Sabaté M, et al. *Prevalence of qnr genes among extended-spectrum b-lactamase-producing enterobacterial isolates in Barcelona, Spain*. J Antimicrob Chemother 2008;61:291-5.
- [28] Saboohi R, Rajaei B, Sepehri Rad N, et al. *Molecular detection and association of qnrA, qnrB, qnrS and bla_{CMY} resistance genes among clinical isolates of Salmonella spp. in Iran*. Adv Microbiol 2014;4:63-8.
- [29] Pakzad I, Ghafourian S, Taherikalani M, et al. *qnr prevalence in extended spectrum b-lactamases (ESBLs) and none-ESBLs producing Escherichia coli isolated from urinary tract infections in central of Iran*. Iran J Basic Med Sci 2011;14:458-64.
- [30] Raei F, Eftekhari F, Feizabadi MM. *Prevalence of quinolone resistance among extended-spectrum β -lactamase producing uropathogenic Klebsiella pneumoniae*. Jundishapur J Microbiol 2014;7:e10887.
- [31] Zamani A, Yousefi Mashouf R, Ebrahimzadeh Namvar AM, et al. *Detection of magA gene in Klebsiella spp. isolated from clinical samples*. Iran J Basic Med Sci 2013;16:173-6.
- [32] Kim HB, Park CH, Kim CJ, et al. *Prevalence of plasmid-mediated quinolone resistance determinants over a 9-year period*. Antimicrob Agents Chemother 2009;53:639-45.
- [33] Wang A, Yang Y, Lu Q, et al. *Presence of qnr gene in Escherichia coli and Klebsiella pneumoniae resistant to ciprofloxacin isolated from pediatric patients in China*. BMC Infectious Diseases 2008;8:1-6.
- [34] Dahmen S, Poirel L, Mansour W, et al. *Prevalence of plasmid-mediated quinolone resistance determinants in Enterobacteriaceae from Tunisia*. Clin Microbiol Infect 2010;16:1019-23.
- [35] Yan Jiang, Zhihui Zhou, Ying Qian, et al. *Plasmid-mediated quinolone resistance determinants qnr and aac(6')-Ib-cr in extended-spectrum b-lactamase-producing Escherichia coli and Klebsiella pneumoniae in China*. J Antimicrob Chemother 2008;61:1003-6.
- [36] Wang M, Sahm DF, Jacoby GA, et al. *Emerging plasmid-mediated quinolone resistance associated with the qnr gene in Klebsiella pneumoniae clinical isolates in the United States*. Antimicrob Agents Chemother 2004;48:1295-9.
- [37] Bouchakour M, Zerouali K, Gros Claude JD, et al. *Plasmid-mediated quinolone resistance in expanded spectrum beta lactamase producing enterobacteriaceae in Morocco*. J Infect Dev Ctries 2010;4:779-803.
- [38] Seyedpour SM, Eftekhari F. *Quinolone susceptibility and detection of qnr and aac(6')-Ib-cr genes in community isolates of Klebsiella pneumoniae*. Jundishapur J Microbiol 2014;7:e11136.
- [39] Wu JJ, Ko WC, Tsai SH, et al. *Prevalence of plasmid-mediated quinolone resistance determinants QnrA, QnrB, and QnrS among clinical isolates of Enterobacter cloacae in a Taiwanese hospital*. Antimicrob Agents Chemother 2007;51:1223-7.
- [40] Robicsek A, Strahilevitz J, Sahm DF, et al. *qnr prevalence in ceftazidime-resistant Enterobacteriaceae isolates from the United States*. Antimicrob Agents Chemother 2006;50:2872-4.

■ Received on January 2, 2015. Accepted on April 1, 2015.

■ Correspondence: Ali Asghar Pahlevan, Cellular and Molecular Research Center, Qazvin University of Medical Sciences, Qazvin, Iran Tel. +98 (28) 33324971- Fax +98(28)33324971 - E-mail: ali_pahlevan@yahoo.com