

A *Clostridium difficile* outbreak in an Italian hospital: the efficacy of the multi-disciplinary and multifaceted approach

A.M. SPAGNOLO^{1,2}, M. SARTINI^{1,2}, A. BATTISTELLA³, B. CASINI⁴, G. LO PINTO³, E. SCHINCA^{1,2}, M.L. CRISTINA^{1,2}
AND HOSPITAL INFECTION CONTROL OPERATING GROUP GALLIERA HOSPITAL³

¹UO SSD Hospital Hygiene, Galliera Hospital, Genoa, Italy; ²Department of Health Sciences, University of Genoa, Italy; ³Galliera Hospital, Genoa, Italy; ⁴Department of Translational Research, New Technologies in Medicine and Surgery, University of Pisa, Italy

Keywords

C. difficile • Outbreak • Infection control practices • Implementation

Summary

Introduction. We described an outbreak of *C. difficile* that occurred in the Internal Medicine department of an Italian hospital and assessed the efficacy of the measures adopted to manage the outbreak.

Methods. The outbreak involved 15 patients and was identified by means of continuous integrated microbiological surveillance, starting with laboratory data (alert organism surveillance). Diarrheal fecal samples from patients with suspected infection by *C. difficile* underwent rapid membrane immuno-enzymatic testing, which detects both the presence of the glutamate dehydrogenase antigen and the presence of the A and B toxins. Extensive microbiological sampling was carried out both before and after sanitation of the environment, in order to assess the efficacy of the sanitation procedure.

Results. The outbreak lasted one and a half month, during which time the Committee for the Prevention of Hospital Infec-

tions ordered the implementation of multiple interventions, which enabled the outbreak to be controlled and the occurrence of new cases to be progressively prevented. The strategies adopted mainly involved patient isolation, reinforcement of proper hand hygiene techniques, antimicrobial stewardship and environmental decontamination by means of chlorine-based products. Moreover, the multifaceted management of the outbreak involved numerous sessions of instruction/training for nursing staff and socio-sanitary operatives during the outbreak. Sampling of environmental surfaces enabled two sites contaminated by *C. difficile* to be identified.

Conclusions. Joint planning of multiple infection control practices, together with effective communication and collaboration between the Hospital Infections Committee and the ward involved proved to be successful in controlling the outbreak.

Introduction

C. difficile is a Gram-positive anaerobic bacterium. Its vegetative cells are capable of forming spores, which confer resistance to heating, drying and chemical agents, including disinfectants. The pathogenic strains of *C. difficile* produce large exotoxin proteins, toxin A (TcdA) and toxin B (TcdB), which constitute the principal virulence factors of the microorganism [1, 2].

Disease caused by *C. difficile* can range in severity from mild diarrhea to fulminant pseudomembranous colitis and, without suitable treatment, toxic megacolon and death [3]. A recent prevalence survey of healthcare-associated infections (HAI) conducted in 183 hospitals determined that *C. difficile* was the most frequently reported infectious agent, being responsible for 12.1% of all HAI [4, 5].

Clostridium difficile has increased in prevalence since 2000, and has caused outbreaks of nosocomial diarrhea worldwide [6]. The main cause of most outbreaks of *Clostridium difficile* infection is NAP1/BI/027: a more virulent ribotype that has been associated with signifi-

cantly higher morbidity and mortality as a result of more severe complications [7]. It is characterized by an *in vitro* overproduction of toxins A and B and by the production of binary toxins [2].

The principal risk factor in *Clostridium difficile* infection (CDI) is antibiotic use, and antibiotics from almost all classes have been associated with infection [7]. Other well-described risk factors are: advanced age, extensive comorbidity, and prolonged hospital stay leading to asymptomatic carriage, recurrent diarrhea, pseudomembranous colitis, or death [7-10].

Patients suffering from *C. difficile* infection shed large amounts of spores that are resistant to disinfectants and regular cleaning procedures, contaminating their surroundings and the hands of nurses, medical staff and others who come into contact with them; hence, contaminated environmental surfaces play a major role in the transmission of *C. difficile* in hospitals [11, 12].

The mortality associated with CDI is high, particularly in older adults with comorbid conditions, severe disease and illness caused by the NAP1 strain of *C. difficile* [13]. Mortality is at least 6% within 3 months of diagnosis and 13% in patients >80 years of age [14].

The economic impact of CDI on the healthcare system is significant, as it doubles the average length of hospitalization and increases the cost of treatment [6, 15]. Nosocomial transmission highlights the importance of rigorous infection control practices for preventing the spread of *C. difficile* [14, 16].

The aims of the present study were to describe an outbreak of *C. difficile* that occurred from 29 December 2015 to 15 February 2016 in the Internal Medicine department of an Italian hospital and to assess the efficacy of the measures adopted to manage the outbreak.

Methods

The outbreak occurred in a nationally renowned, highly specialized hospital in northern Italy, organized in accordance with treatment intensity. The facility is composed of separate pavilions with a total of 431 beds. The ward directly involved was female internal medicine, which has 26 beds.

Hospital infection cases were defined as patients with positive toxin assays > 48 hours after hospital admission. The outbreak, which involved 15 patients from 29 December 2015 to 15 February 2016, was identified by means of continuous integrated microbiological surveillance, starting with laboratory data (alert organism surveillance). Following laboratory identification of an epidemiologically important microorganism, the dedicated software of the surveillance system automatically e-mails the data to all the members of the Hospital Infections Committee (made up of members of the hospital's healthcare administration, physicians, microbiologists, infectious disease specialists, epidemiologists), who then implement the interventions deemed necessary, with particular regard to the application of isolation measures. A validated report is simultaneously sent through the laboratory information system to the hospital facility involved.

For patients with a diagnosis of *Clostridium difficile*, information on age, history of hospitalizations, antibiotic treatments, duration of hospitalization and outcome were collected.

MICROBIOLOGICAL ANALYSIS

Diarrheal fecal samples from patients with suspected infection by *C. difficile* underwent rapid membrane immuno-enzymatic testing by means of the TECHLAB C. diff Chek Quick Complete® (Alere™) kit, which detects both the presence of the glutamate dehydrogenase (GDH) antigen, as a means of screening for *C. difficile*, and the presence of the A and B toxins.

ENVIRONMENTAL INVESTIGATION

Extensive microbiological sampling was carried out both before and after sanitation of the environment, in order to assess the efficacy of the sanitation procedure. Sampling was carried out at 14 sites of high-frequency contact; the sampling points were selected in accordance with the checklist of the CDCs reported in the

APIC guidelines "Guide to Preventing *Clostridium difficile* Infections" [17], which specifies the critical points to be examined in the event of an outbreak. Monitoring therefore included critical surfaces in proximity to the patient's bed (e.g. personal light switch and call button) and other surfaces at high risk of contact with hospital personnel (e.g. medicine trolley, light switch, curtains between the beds, etc) or patients.

In accordance with the methods of Best et al. [18] and Ali et al. [19], specimens were taken by using 25-cm² sponge swabs pre-moistened with neutralizing solution (Medical Wire & Equipment, England). The swabs were then placed aseptically into sterile Stomacher bags containing 50 ml of Ringer solution (Oxoid) and homogenized manually by vigorously massaging the bag between the fingertips for 1 min. Liquid from the bag was passed through a 0.45-µm filter (Millipore), which was then placed aseptically onto Brazier's *Clostridium difficile* selective agar (Oxoid). Plates were then incubated at 37°C under anaerobic conditions for 48 h prior to reading.

C. difficile was initially identified on the basis of the macroscopic appearance of colonies and microscopic characteristics, and confirmed to be *C. difficile* by means of latex agglutination testing (Oxoid *C. difficile* Test kit).

Results

DESCRIPTION OF THE OUTBREAK

Following the analysis of patients' records, a possible index case was identified: an 86-year-old woman hospitalized on 16 December 2015 in the ward where the outbreak originated. This patient had already been admitted to the same hospital in the previous month (geriatric ward) for bronchopneumopathy.

On 12 December she was taken to the Emergency Department with bruising to the pelvis after a fall at home. A bilateral pleural effusion and respiratory insufficiency were diagnosed. She was therefore hospitalized in the Sub-intensive Care Unit and, after being stabilized, was transferred to the Internal Medicine Department three days later.


Table I reports the characteristics of the patients involved in the outbreak. Their mean age was 82.13 years (range 70-90 years), the mean Charlson index was 7 (range 4-13) and the mean duration of hospitalization before the first isolation of *C. difficile* was 16 days (range 4-34 days). With regard to outcome, 5 patients died (4 attributable to *C. difficile*), 6 were transferred to other healthcare facilities and/or wards, and 4 were discharged.

In 86.67% of cases, the *C. difficile* strain responsible for the infection produced both toxin A and toxin B; in the remaining cases, weak positivity to the immuno-enzymatic test was recorded. The patients were treated with metronidazole and, in the event of failure, vancomycin. Figure 1 describes distribution of *Clostridium difficile* infected patients as a function of time

Tab. I. Characteristics of patients involved in the outbreak

Patient	<i>Clostridium difficile</i> toxins	Days of hospitalization	Age	Charlson index	Bed	Outcome
1*	A and B	16	85	7	23	Transferred
2	A and B	11	80	7	24	Transferred
3	A and B	9	70	12	6	Died
4	A and B	27	88	6	5	Died
5	A and B	26	79	4	8	Discharged
6	Weak positivity	13	84	6	9	Discharged
7	A and B	16	75	7	1	Died
8	A and B	12	85	5	10	Discharged
9	A and B	15	90	8	7	Died
10	A and B	4	84	5	1	Discharged
11	A and B	34	80	6	14	Died
12	A and B	16	77	4	23	Transferred
13	Weak positivity	12	80	7	10	Transferred
14	A and B	17	88	6	6	Transferred
15	A and B	12	87	13	26	Transferred

*Suspected index case




INFECTION CONTROL PRACTICES DURING THE OUTBREAK

Figure 2 shows the epidemic curve of the outbreak from 29 December 2015 to 15 February 2016. The timescale of the interventions implemented by the Committee for the Prevention of Hospital Infections is also indicated. From the moment when the first two cases of infection were diagnosed an antimicrobial stewardship program and the following interventions were implemented:

Intervention 1: from 29 December 2015:

- Specification of the measures to be taken in order to contain risk of infection by *Clostridium difficile*, considering all patients to be potentially infected; written instructions delivered to all healthcare personnel involved.
- Testing for *C. difficile* toxins in all symptomatic patients.
- Isolation in cohorts of infected patients; assistance to cohorts (dedicated operators); use of dedicated small devices (e.g. oximeter, hemoglucotest device, etc) for infected patients.

- *Ad hoc* environmental sanitation for *Clostridium difficile* in the entire department (with 20% concentrations of chlorine-based detergent), including decontamination of telephones and computer keyboards and screens (ready-to-use sodium hypochlorite solution). In order to facilitate adequate daily sanitation, bedside tables were kept clear of all but indispensable objects (bottle of water and glass).
- Checking to ensure that healthcare personnel complied with hand hygiene protocols. In addition, the hands of all non-self-sufficient patients were washed more frequently and self-sufficient patients were instructed on how to wash their hands properly.
- Checking to ensure that gloves were used properly and were changed after assisting each individual patient, and that hands were washed immediately after the removal of gloves.
- Operators involved in direct assistance were instructed to change their overalls daily and were encouraged to use microfiber overalls, which are more protective of the hygiene of infected patients, and disposable nonwoven gowns.
- Staff were forbidden to use personal mobile phones while assisting infected patients.

Fig. 2. Epidemic curve of the *C. difficile* outbreak and the timescale of the interventions implemented.

- Correct patient hygiene practices were emphasized; soiled underwear was placed in an impermeable bag labeled with the patient's name, which was then placed in a dedicated container inside the room/cubicle. If a patient lift was used, the sling cover was changed for each patient and sent for disinfection as if it were certainly infected; the same approach was adopted towards minor aids, for which disposable protective covers were also used.
- The number of visitors was reduced, and a specific information leaflet concerning the behavior of visitors to infected patients was distributed; this provided instructions on hand washing and interpersonal contact.

The day after implementation of intervention 1, another two cases of infection were discovered. Following a meeting to update and instruct nursing staff and social/healthcare workers, the second phase of intervention was implemented.

Intervention 2: from 30 December 2015:

- Simulation of donning and removing personal protection devices (PPD).
- Reiteration of procedures for the proper sanitation of stands for i.v. drips, commode chairs, infusion pumps, PCs and telephones.
- Meals served in heat-sealed containers for all patients.

- Checking of proper isolation of infected patients (e.g. collocation of the patient, supply of hand-washing requisites, availability of disposable overalls, materials and dedicated devices, etc.); this revealed the need to supply some types of medical devices for dedicated use (e.g. stethoscope and sphygmomanometer).
- Urgent processing of fecal samples for culture tests; prompt telephone communication of positive reports to the expert consultants of the Committee for the Prevention of Hospital Infections, for immediate application of the necessary measures.
- Periodic checks on compliance with the measures recommended.

Intervention 3: from 5 January 2016

- Ward staff increased on both day shifts and night shifts.

Intervention 4: 8 January 2016

- Review of cases following the administration of antibiotic treatment and implementation of the control measures; assessment of the need to institute further briefings/training for medical and nursing staff.

Intervention 5: from 12 January 2016

- Structural, logistical and organizational segregation of infected patients (left side of the ward) from unin-

fecting patients (right side), and consequent reorganization of the activities of sanitation and assistance.

- Direct observation to ensure proper implementation of the measures to contain the risk of infection, and institution of “on the job” staff training with regard to: donning and removal of personal protection devices; the hygiene of infected patients, with particular regard to the hands; decontamination of the patient-unit; use of personalized devices for each patient; decontamination of the environment, materials and medical devices; functional isolation of cohorts; institution of a differential pathway from “clean” to “dirty”; proper collection and conservation of fecal samples prior to analysis; application of medication to CVC with maintenance of asepsis in infected patients; healthcare education of visitors, with simulation of hand hygiene.

From this date onwards, thanks to the set of control measures adopted, the number of cases of infection progressively diminished, and the last two cases recorded on 8 February 2016 were resolved.

ENVIRONMENTAL MONITORING

The microbiological results of environmental monitoring conducted on 15 January 2016 revealed contamination by *Clostridium difficile* on the curtain separating two beds that had been occupied by patients involved in the outbreak (beds 23 and 24) and on the call button of bed 24. The curtain was promptly removed and disposed of, and the entire environment was thoroughly disinfected. Subsequent monitorings, carried out after environmental sanitation, revealed no contamination by *C. difficile*.

Discussion

Several reports suggest that the incidence and severity of *C. difficile* infection have been increasing in recent years across the United States, Canada and Europe. Recent data from 28 community hospitals in the southern United States suggest that *C. difficile* has replaced methicillin-resistant *Staphylococcus aureus* as the most common cause of healthcare-associated infection [20, 21]. The burden of healthcare-associated CDIs in acute-care hospitals in the EU/EEA has been estimated at 123,997 cases annually. In the ECDC point prevalence survey of healthcare-associated infections in European acute-care hospitals 2011-2012, *C. difficile* was the 8th most frequently detected microorganism among HAIs [22].

In the present study, we documented the occurrence of 15 cases of *C. difficile* infection in an internal medicine department in an Italian hospital. During the outbreak the Committee for the Prevention of Hospital Infections ordered the implementation of multiple interventions, which enabled the outbreak to be controlled and the occurrence of new cases to be progressively prevented.

The outbreak described in this paper started and finished in a single ward, involved a relatively small number of patients, and lasted one and a half month. Wong-Mc-

Clure et al. [23] described an outbreak due to *C. difficile* that involved three wards and 389 patients, and which lasted for several months. More recently, van Beurden et al. [6] described an outbreak that involved 19 wards and 72 patients, and which lasted for a year.

As pointed out by several studies, there may not be a single method that is effective in minimizing exposure to *C. difficile*, and a multifaceted approach is usually required [24]. Indeed, the management of CDI in hospitals requires just such a multidisciplinary approach, which begins with infection prevention. A previous study by Weiss et al. [25] showed that a multi-pronged intervention strategy is most effective in reducing the rate of healthcare CDI.

Strategies for the prevention and control of *C. difficile* infections are aimed at promptly identifying, isolating and efficaciously treating patients affected by CDI (in order to reduce the dissemination of spores and prevent secondary cases) and at minimizing preventable risk factors through the implementation of protocols of behavior, environmental sanitation and antibiotic stewardship [26].

In accordance with this approach, the strategies adopted for the control of the *Clostridium difficile* outbreak described here mainly involved patient isolation, reinforcement of proper hand hygiene techniques, antimicrobial stewardship and environmental decontamination by means of chlorine-based products.

Indeed, the presence of other patients with infection, hand carriage on the part of healthcare personnel and contaminated environmental surfaces are considered to be major factors in the transmission of pathogens in hospitals [27-29], including *C. difficile*.

When there is an infected patient in hospital, the hospital environment is contaminated by spores within a few hours of the onset of diarrhea; other patients may therefore be infected and the patient himself/herself may be reinfected. Moreover, *C. difficile* spores are highly resistant to many commonly used disinfectants and may persist for months in hospital environments [30].

Environmental contamination with *C. difficile* spores occurs at as many as 34-58% of sites, despite cleaning, with surfaces of fomites being most frequently contaminated [18].

Frequently touched surfaces in near patient areas are rapidly contaminated by the microorganisms disseminated by the infected patient occupying the room, and may remain contaminated for extended periods of time [31]. Consequently, *C. difficile* can be found on hospital floors, on bedrails, windowsills, commodes, toilets, call buttons, blood pressure cuffs, electronic thermometers, bedsheets and anything that comes into contact with contaminated hands [32]. Thus, thorough disinfection of the contaminated hospital environment is essential in order to prevent the transmission of this nosocomial pathogen, and the choice of hospital decontamination protocols can markedly affect the prevalence and environmental distribution of *C. difficile* contamination [33]. The scientific evidence supports the use of detergents containing chlorine (at least 1000 ppm of active chlo-

rine) in endemic situations or during epidemic outbreaks [1]. A study by Fawley et al. [33] compared the efficacy of five different cleaning agents against epidemic and non-epidemic *C. difficile* strains. They found that only chlorine-based germicides were able to inactivate *C. difficile* spores.

Contamination of the hands of healthcare staff and patients with *C. difficile* is a major route of transmission of the infection, and there is a close correlation between hand contamination and the degree of environmental contamination. For this reason, proper hand hygiene is crucial to preventing the transmission of *C. difficile* in the hospital setting [32].

During the outbreak described in this paper, various interventions were undertaken in order to ensure adherence to hand hygiene protocols on the part of healthcare staff and patients; those visiting infected patients were also taught to wash their hands and to limit contact only to the patient being visited. Indeed, checking the staff's compliance with hand hygiene has been deemed a more effective strategy than microbiological testing of the hands by means of sampling.

A recommendation common to many guidelines on the prevention and control of healthcare-related infections concerns the training of healthcare personnel, visitors, caregivers and patients themselves. The multifaceted management of the outbreak described here involved numerous sessions of instruction/training for nursing staff and socio-sanitary operatives during the course of the epidemic. By modifying risk behaviors, these interventions certainly helped to control the outbreak.

Equally important was environmental monitoring. Limited to times of outbreak, rather than being part of routine practice, this can provide a valuable estimate of the level of contamination on surfaces such as walls, work surfaces, floors and equipment [34] and is currently recommended by the Centers for Disease Control and Prevention [35]. In the present case, sampling of environmental surfaces enabled two sites contaminated by *C. difficile* to be identified, one of which was a soft plastic-coated curtain separating two beds that had previously been occupied by infected patients. As this curtain would have been very difficult to disinfect, it was removed and disposed of immediately after the detection of contamination; this measure may well have enabled an environmental reservoir of the microorganism to be eliminated, a hypothesis that is also supported by the trend in the epidemic curve after the implementation of environmental monitoring.

In conclusion, joint planning of multiple infection control practices, together with effective communication and collaboration between the Hospital Infections Committee and the ward involved proved to be successful in controlling the outbreak.

Acknowledgements

All authors declare that there is no conflict of interest.

Authors' contributions

MLC conceived and designed the study. AB collected data. BC performed the data quality control. ES performed environmental controls. MS validating and analysing the data. MLC and AMS wrote the paper. GLP revised the manuscript. All Authors revised the manuscript and gave their contribution to improve the paper. All authors read and approved the final manuscript.

References

- [1] Cristina ML, Spagnolo AM, Sartini M, Panatto D, Perdelli F. Clostridium difficile infections: An emerging problem in healthcare facilities. *Rev Med Microbiol* 2012;23:67-75.
- [2] Jia H, Du P, Yang H, Zhang Y, Wang J, Zhang W, Han G, Han N, Yao Z, Wang H, Zhang J, Wang Z, Ding Q, Qiang Y, Barbut F, Gao GF, Cao Y, Cheng Y, Chen C. Nosocomial transmission of Clostridium difficile ribotype 027 in a Chinese hospital, 2012-2014, traced by whole genome sequencing. *BMC Genomics* 2016;17:405.
- [3] Secore S, Wang S, Dougherty J, Xie J, Miezejewski M, Rustandi RR, Horton M, Xoconostle R, Wang B, Lancaster C, Kristopeit A, Wang SC, Christanti S, Vitelli S, Gentile MP, Goerke A, Skinner J, Strable E, Thiriot DS, Bodmer JL, Heinrichs JH. Development of a novel vaccine containing binary toxin for the prevention of clostridium difficile disease with enhanced efficacy against NAP1 Strains. *PLoS One* 2017;12(1):e0170640.
- [4] Magill SS, Edwards JR, Bamberg W, Beldavs ZG, Dumyati G, Kainer MA, Lynfield R, Maloney M, McAllister-Holod L, Nade J, Ray SM, Thompson DL, Wilson LE, Fridkin SK; Emerging Infections Program Healthcare-Associated Infections and Antimicrobial Use Prevalence Survey Team. Multistate point-prevalence survey of health care-associated infections. *N Engl J Med* 2014;370:1198-208.
- [5] Morfin-Otero R, Garza-Gonzalez E, Aguirre-Diaz SA, Escobedo-Sanchez R, Esparza-Ahumada S, Perez-Gomez HR, Petersen-Morfin S, Gonzalez-Diaz E, Martinez-Melendez A, Rodriguez-Noriega E; Hospital Civil de Guadalajara, Fray Antonio Alcalde Clostridium difficile Team. Clostridium difficile outbreak caused by NAP1/BI/027 strain and non-027 strains in a Mexican hospital. *Braz J Infect Dis* 2016;20:8-13.
- [6] van Beurden YH, Bomers MK, van der Werff SD, Pompe EA, Spiering S, Vandenbroucke-Grauls CM, Mulder CJ. Cost analysis of an outbreak of Clostridium difficile infection ribotype 027 in a Dutch tertiary care centre. *J Hosp Infect* 2017;95:421-5.
- [7] van Beurden YH, Dekkers OM, Bomers MK, Kaiser AM, van Houdt R, Knetsch CW, Girbes AR, Mulder CJ, Vandenbroucke-Grauls CM. An outbreak of Clostridium difficile ribotype 027 associated with length of stay in the intensive care unit and use of selective decontamination of the digestive tract: A Case Control Study. *PLoS One* 2016;11:e0160778.
- [8] Kumar N, Miyajima F, He M, Roberts P, Swale A, Ellison L, Pickard D, Smith G, Molyneux R, Dougan G, Parkhill J, Wren BW, Parry CM, Pirmohamed M, Lawley TD. Genome-based Infection tracking reveals dynamics of Clostridium difficile transmission and disease recurrence. *Clin Infect Dis* 2016;62:746-52.
- [9] Visconti V, Brunetti G, Cuomo MR, Giordano A, Raponi G. Nosocomial-acquired and community-onset Clostridium difficile infection at an academic hospital in Italy: Epidemiology, recurrences and toxin genes distribution. *J Infect Chemother* 2017; doi: 10.1016/j.jiac.2017.08.002.
- [10] Vonberg RP, Kuijper EJ, Wilcox MH, Barbut F, Tüll P, Gastmeier P, European C difficile-Infection Control Group; European Centre for Disease Prevention and Control (ECDC), van den Broek PJ, Colville A, Coignard B, Daha T, Debast S, Duerden BI, van den Hof S, van der Kooij T, Maarleveld HJ, Nagy E,

- Notermans DW, O'Driscoll J, Patel B, Stone S, Wiuff C. Infection control measures to limit the spread of *Clostridium difficile*. *Clin Microbiol Infect* 2008;14:2-20.
- [11] Morales L, Rodríguez C, Gamboa-Coronado MD. Molecular detection of *Clostridium difficile* on inert surfaces from a Costa Rican hospital during and after an outbreak. *Am J Infect Control* 2016;44:1517-9.
- [12] Cristina ML, Spagnolo AM, Orlando P, Perdelli F. The role of the environment in the spread of emerging pathogens in at-risk hospital wards. *Rev Med Microbiol* 2013;104-12.
- [13] Evans CT, Safdar N. Current trends in the epidemiology and outcomes of *Clostridium difficile* infection. *Clin Infect Dis* 2015;60:S66-71.
- [14] Jullian-Desayes I, Landelle C, Mallaret MR, Brun-Buisson C, Barbut F. *Clostridium difficile* contamination of health care workers' hands and its potential contribution to the spread of infection: Review of the literature. *Am J Infect Control* 2017;45:51-8.
- [15] Bouza E. Consequences of *Clostridium difficile* infection: understanding the healthcare burden. *Clin Microbiol Infect* 2012;18:5-12.
- [16] Planche T, Aghaizu A, Holliman R, Riley P, Poloniecki J, Breathnach A, Krishna S. Diagnosis of *Clostridium difficile* infection by toxin detection kits: a systematic review. *Lancet Infect Dis* 2008;8:777-84.
- [17] APIC Implementation Guide 2013. Guide to Preventing *Clostridium difficile* Infections. 2013 http://apic.org/Resource/_EliminationGuideForm/59397fc6-3f90-43d1-9325-e8be75d86888/File/2013CDiffFinal.pdf. Accessed 30 Dec 2015.
- [18] Best EL, Fawley WN, Parnell P, Wilcox MH. The potential for airborne dispersal of *Clostridium difficile* from symptomatic patients. *Clin Infect Dis* 2010;50:1450-7.
- [19] Ali S, Muzslay M, Wilson P. A novel quantitative sampling technique for detection and monitoring of *Clostridium difficile* contamination in the clinical environment. *J Clin Microbiol* 2015;53:2570-4.
- [20] Lessa FC, Gould CV, McDonald LC. Current status of *Clostridium difficile* infection epidemiology. *Clin Infect Dis* 2012;55:S65-70.
- [21] Miller BA, Chen LF, Sexton DJ, Anderson DJ. Comparison of the burdens of hospital-onset, healthcare facility-associated *Clostridium difficile* Infection and of healthcare-associated infection due to methicillin-resistant *Staphylococcus aureus* in community hospitals. *Infect Control Hosp Epidemiol* 2011;32:387-90.
- [22] ECDC. *Clostridium difficile* infections. 2016. http://antibiotic.ecdc.europa.eu/en/healthtopics/Healthcare-associated_infections/Clostridium-difficile-infections/Pages/Clostridium-difficile-infections.aspx Accessed 3 Apr 2017.
- [23] Wong-McClure RA, Ramírez-Salas E, Mora-Brenes N, Agüero-Sandí L, Morera-Sigler M, Badilla-Vargas X, Hernández-de Merzerville M, O'Shea M, Bryce E. Long term effect of infection control practices and associated factors during a major *Clostridium difficile* outbreak in Costa Rica. *J Infect Dev Ctries* 2013;7:914-21.
- [24] Cohen SH, Gerding DN, Johnson S, Kelly CP, Loo VG, McDonald LC, Pepin J, Wilcox MH; Society for Healthcare Epidemiology of America; Infectious Diseases Society of America. Clinical practice guidelines for *Clostridium difficile* infection in adults: 2010 update by the society for healthcare epidemiology of America (SHEA) and the infectious diseases society of America (IDSA). *Infect Control Hosp Epidemiol* 2010;31:431-55.
- [25] Weiss K, Boisvert A, Chagnon M, Duchesne C, Habash S, Lepage Y, Letourneau J, Raty J, Savoie M. Multipronged intervention strategy to control an outbreak of *Clostridium difficile* infection (CDI) and its impact on the rates of CDI from 2002 to 2007. *Infect Control Hosp Epidemiol* 2009;30:156-62.
- [26] SIMPIOS. Prevenzione e controllo delle infezioni da *Clostridium difficile*. 2009. <http://www.simpio.it/public/ufiles/Prevenzione%20e%20controllo%20delle%20infezioni%20da%20Clostridium%20Difficile.pdf> Accessed 30 Dec 2015.
- [27] Perdelli F, Dallera M, Cristina ML, Sartini M, Ottria G, Spagnolo AM, Orlando P. A new microbiological problem in intensive care units: environmental contamination by MRSA with reduced susceptibility to glycopeptides. *Int J Hyg Environ Health* 2008;211:213-8.
- [28] Orlando P, Cristina ML, Dallera M, Ottria G, Vitale A, Badolati G. Surface disinfection: evaluation of the efficacy of a nebulization system spraying hydrogen peroxide. *J Prev Med Hyg* 2008;49:116-9.
- [29] Spagnolo AM, Orlando P, Panatto D, Amicizia D, Perdelli F, Cristina ML. *Staphylococcus aureus* with reduced susceptibility to vancomycin in healthcare settings. *J Prev Med Hyg* 2014;55:137-44.
- [30] Kuijper EJ, Coignard B, Tüll P. Emergence of *Clostridium difficile*-associated disease in North America and Europe. *Clin Microbiol Infect* 2006;12:2-18.
- [31] Ali S, Muzslay M, Bruce M, Jeanes A, Moore G, Wilson AP. Efficacy of two hydrogen peroxide vapour aerial decontamination systems for enhanced disinfection of methicillin-resistant *Staphylococcus aureus*, *Klebsiella pneumoniae* and *Clostridium difficile* in single isolation rooms. *J Hosp Infect* 2016;93:70-7.
- [32] Martinez FJ, Leffler DA, Kelly CP. *Clostridium difficile* outbreaks: prevention and treatment strategies. *Risk Manag Healthc Policy* 2012;5:55-64.
- [33] Fawley WN, Underwood S, Freeman J, Baines SD, Saxton K, Stephenson K, Owens RC Jr, Wilcox MH. Efficacy of hospital cleaning agents and germicides against epidemic *Clostridium difficile* strains. *Infect Control Hosp Epidemiol* 2007;28:920-5.
- [34] Ottria G, Dallera M, Aresu O, Manniello MA, Parodi B, Spagnolo AM, Cristina ML. Environmental monitoring programme in the cell therapy facility of a research centre: preliminary investigation. *J Prev Med Hyg* 2010;51:133-8.
- [35] Sehulster L, Chinn R, CDC, HICPAC. Guidelines for environmental infection control in health-care facilities. Recommendations of CDC and the Healthcare Infection Control Practices Advisory Committee (HICPAC). *MMWR Recomm Rep* 2003;52:1-42.

■ Received on May 18, 2018. Accepted on May 30, 2018.

■ Correspondence: Marina Sartini, Department of Health Sciences, University of Genoa, via Pastore 1, 16132 Genova, Italy - Tel. +39 010 353 8447 - Fax +39 010 353 8216 - E-mail: sartini@unige.it